

Anomalous energy transport in FPU- β chain

A. Mellet* and S. Merino-Aceituno†

November 20, 2014

Abstract

This paper is devoted to the derivation of a macroscopic diffusion equation (Fourier's law) describing the transport of energy in an anharmonic chain. More precisely, we study here the so-called FPU- β chain, which is a very simple model for a one-dimensional crystal in which atoms are coupled to their nearest neighbors by a harmonic potential, weakly perturbed by a quartic potential. The starting point of our mathematical analysis is a kinetic equation: Lattice vibrations, responsible for heat transport, are modeled by an interacting gas of phonons whose evolution is described by the Boltzmann Phonon Equation. Our main result is the derivation of an anomalous diffusion equation for the temperature.

1 Introduction

In this paper, we investigate some aspects of the transport of energy in one dimensional chains of oscillators. The goal is to derive Fourier's law, which is at the core of the heat equation and states that the heat flux \vec{j} behaves as

$$\vec{j} = -\kappa \nabla_x T \quad (1)$$

where T is the temperature and κ is a positive constant that may depend on the temperature itself. This law has been observed experimentally, but there is no rigorous mathematical justification describing how it arises from the atomic laws of the solid (see [10] for a review of this very challenging problem).

At the microscopic level, solids can be modeled as lattices, where each node represents an atom. For insulating crystals, where heat is transported

*University of Maryland, College Park 20742, USA.

†University of Cambridge, Cambridge CB3 0WA, UK

by lattice vibrations (see [13]), one possible approach to derive Fourier's law relies on the introduction of the Boltzmann phonon equation, a kinetic equation that can play the role of an intermediate step between the microscopic atomic level and the macroscopic scale. It is this approach, first suggested by Peierls [20], that we try to make rigorous in this paper in a very particular setting.

The particular framework we are considering was made popular by a famous numerical experiment performed by Fermi, Pasta and Ulam in the 1950's at Los Alamos National Laboratories. The goal of their experiment was to investigate numerically the dynamic (and relaxation toward equilibrium) of the simplest model for a crystal: a chain of oscillators coupled to their nearest neighbors by non-linear forces described by an Hamiltonian of the form

$$H = \frac{1}{2} \sum_{i \in \mathbb{Z}} [p_i^2 + V(q_{i+1} - q_i)].$$

When V is purely harmonic, the system has quasi-periodic solutions and does not relax to an equilibrium (see [9]). Fermi, Pasta and Ulam thus considered the next two simplest cases by adding a cubic potential $V(r) = r^2 + \alpha r^3$ (this model is now referred to as the FPU- α chain) or a quartic potential $V(r) = r^2 + \beta r^4$ (the FPU- β chain).

These models have been widely studied since that original experiment (see Lepri, Livi, Politi [14] for a recent review of the work devoted to these models). Our goal in this paper is to derive Fourier's law for the FPU- β chain (we will see later why we do not consider the FPU- α chain). To achieve this, we rely on an idea of Peierls [20], who describes lattice vibrations, responsible for heat transport, as an interacting gas of phonons whose density distribution function (denoted W below) solves a Boltzmann phonon equation (also known as Peierls equation in this context). The mathematical derivation of this Boltzmann phonon equation starting from the microscopic equations for the motion of the atoms (Hamiltonian dynamic) has been made more rigorous recently by H. Spohn in [24]. We will thus not focus on this step, though we will spend some time in this paper discussing the results of [24]. Our focus instead will be on the rigorous derivation of Fourier's law from the Boltzmann phonon equation. The most remarkable aspect of our result is that we will not recover (1), but instead a non-local Fourier law corresponding to an anomalous diffusion equation (in place of the usual heat equation). This was not unexpected, since anomalous heat diffusion phenomena in the FPU- β framework has actually been observed numerically in dimension one and two (while normal diffusion is observed in the three

dimensional case), see in particular [22], [13], [14], and also [3] for a study at the level of the kinetic equation. In fact, by using Peierls equation, it has previously been proved that the energy current correlation has a slow decay in time as $t^{-3/5}$ indicating anomalous diffusive behavior (see [21, 15]).

Let us now describe our main result. As mentioned above, the starting point of our analysis is the Boltzmann phonon equation given by:

$$\partial_t W + \omega'(k)\partial_x W = C(W)$$

where the unknown $W(t, x, k)$ is a function of the time $t \geq 0$, the position $x \in \mathbb{R}$ and the wave vector $k \in \mathbb{T} := \mathbb{R}/\mathbb{Z}$. This function is introduced in [24] as the Wigner transform of the displacement field of the atoms, but it can be interpreted as a density distribution function for a gas of interacting phonons (describing the chain vibrations). The function $\omega(k)$ is the dispersion relation for the lattice and the operator C describes the interactions between the phonons.

We will discuss in Sections 2 and 3 the particular form of ω and C corresponding to our microscopic models. For the FPU- β chain, the operator C will be the so-called *four phonon collision operator*, which is an integral operator of Boltzmann type but cubic instead of quadratic (see (19)).

As explained above, our goal is to derive a macroscopic equation for the temperature. This is, at least in spirit, similar to the derivation of Navier-Stokes equations from the Boltzmann equation for diluted gas (see [4] and references therein). We will consider a perturbation of a thermodynamical equilibrium $\bar{W}(k) = \frac{\bar{T}}{\omega(k)}$ (note that the temperature is classically defined by the relation $E = k_B T$ where $E = \int_{\mathbb{T}} \omega(k) W(k) dk$ and k_B denotes Boltzmann's constant - here, we choose temperature units so that $k_B = 1$):

$$W^\varepsilon(t, x, k) = \bar{W}(k)(1 + \varepsilon f^\varepsilon(t, x, k)).$$

The function f^ε then solves

$$\partial_t f^\varepsilon + \omega'(k)\partial_x f^\varepsilon = L(f^\varepsilon) + \mathcal{O}(\varepsilon)$$

where L is the linearized operator

$$L(f) = \frac{1}{\bar{W}} DC(\bar{W})(\bar{W}f).$$

As usual a macroscopic equation is derived after an appropriate rescaling of the time and space variable. More precisely, we will show (see Theorem 4.1) that the solution of

$$\varepsilon^{\frac{8}{5}} \partial_t f^\varepsilon + \varepsilon \omega'(k)\partial_x f^\varepsilon = L(f^\varepsilon)$$

converges to a function $T(t, x)$ solution of

$$\partial_t T + \frac{\kappa}{\bar{T}^{6/5}} (-\Delta)^{4/5} T = 0$$

thus giving the anomalous Fourier's law (of order 3/5)

$$\vec{j} = -\kappa(\bar{T}) \nabla (-\Delta)^{-1/5} T.$$

The derivation of such a fractional diffusion equation from a kinetic equation is now classical (see [17], [16], [8], [1], [12]). As in previous results (see in particular [8]), the order of the limiting diffusion process is determined by the degeneracy of the collision frequency of the operator L . Our work is thus greatly indebted to the work of J. Lukkarinen and H. Spohn [15] who carefully study the properties of the operator L and show in particular that the collision frequency behaves as $|k|^{5/3}$ as $k \rightarrow 0$.

The main novelty here, compared with the results mentioned above, is the fact that the kernel of the collision operator L is 2 dimensional. The reason for that will be discussed in the next sections and it appears to be a mathematical artifact rather than being related to some physical phenomenon. It does, however, indicate some weakness in the mixing properties of the collision process (this will be even more obvious for the FPU- α chain, for which the collision operator vanishes altogether). And while the macroscopic behavior of f^ε is completely determined by the function $T(t, x)$, the other component of the projection of f^ε onto the kernel of L will play a role in reducing the value of the diffusion coefficient κ .

We point out that we will not attempt here to derive a nonlinear Fourier law by working with the nonlinear operator C (rather than the linearized operator L). Such a derivation is developed in [11] by Bricmont and Kupiainen, but under assumptions that ensure that regular diffusion, rather than anomalous diffusion, takes place (non degeneracy of the collision frequency).

We conclude this introduction by mentioning that diffusive and superdiffusive heat transport has also been derived for FPU-type chains in a different mathematical setting using a probabilistic approach: in this setting the hamiltonian dynamics of the microscopic system are considered to have only an harmonic potential and the dynamics are perturbed by a stochastic noise conserving momentum and energy (see [5], [6], [7] and the review paper [19].)

This paper is organized as follows: In the next section, we describe the original problem (chains of coupled harmonic oscillators) and its relation to the Boltzmann phonon equation. We then introduce the collision operators

C that appears in the context of FPU chains. In that section, we will see in particular that this kinetic description cannot be used to study the FPU- α chain because the collision operator C vanishes in that case. This section is mostly based on the paper of H. Spohn [24].

In Section 3, we investigate the properties of the four phonon collision operators, appearing in the context of the FPU- β chain as well as its linearization around an equilibrium (this section is largely based on the work of J. Lukkarinen and H. Spohn [15]). The main result of our paper is finally stated in Section 4 and its proof is divided between Sections 5 and 6.

Acknowledgement

This material is based upon work supported by the Kinetic Research Network (KI-Net) under the NSF Grant No. RNMS #1107444.

A.M. is partially supported by NSF Grant DMS-1201426.

S.M thanks the University of Maryland and CSCAMM (Center for Scientific Computation and Mathematical Modeling) for their hospitality; the Cambridge Philosophical Society and Lucy Cavendish College (University of Cambridge) for their financial support. Thanks to Clément Mouhot from the University of Cambridge for his help and support. Thanks to Herbert Spohn from Technische Universität München for useful discussions. S.M is supported by the UK Engineering and Physical Sciences Research Council (EPSRC) grant EP/H023348/1 for the University of Cambridge Centre for Doctoral Training, the Cambridge Centre for Analysis.

2 Crystal vibrations: A kinetic description

In this section, we recall the results from the paper of H. Spohn [24] that are relevant to our present study. Our goal is to detail the relation between the Boltzmann phonon equation that we are considering in this paper and the microscopic models. At the microscopic level, we consider an infinite lattice \mathbb{Z}^n describing the equilibrium positions of the atoms of a crystal (we briefly introduce the model in general dimension, though starting in the next section, we will focus solely on the one-dimensional case). The deviation of the atom $i \in \mathbb{Z}^n$ from its equilibrium position is denoted by q_i , and the conjugate momentum variable is denoted by p_i . We consider the dynamics

associated to the Hamiltonian

$$H(q, p) = \frac{1}{2} \sum_{i \in \mathbb{Z}} p_i^2 + V_h(q) + \sqrt{\lambda} V(q)$$

where V_h is a harmonic potential (quadratic) and $\sqrt{\lambda} V$ is a small anharmonic perturbation (the kinetic equation is obtained in the limit $\lambda \rightarrow 0$). The general form of the harmonic potential is

$$V_h(q) = \frac{1}{2} \sum_{i, j \in \mathbb{Z}^n} \alpha(i - j) q_i q_j + \frac{\omega_0^2}{2} \sum_{i \in \mathbb{Z}^n} q_i^2, \quad (2)$$

while V is typically a cubic or quartic potential of the form

$$V(q) = \sum_{i \in \mathbb{Z}^n} \gamma(q_i) \quad \text{or} \quad V(q) = \sum_{\substack{i, j \in \mathbb{Z}^n \\ |i - j| = 1}} \gamma(q_j - q_i).$$

In order to understand how energy is being transported by the vibration of the atoms in the lattice, we will replace this very large system of ODE by a kinetic equation (the so-called Boltzmann phonon equation) whose unknown $W(x, k, t)$ will be interpreted as a density distribution function for a gas of interacting phonons. The idea of describing the lattice vibrations by interacting phonons, whose evolution would be described by a Boltzmann type equation first appeared in a paper of Peierls [20]. This derivation was made more rigorous by H. Spohn [24] using Wigner transforms and asymptotic analysis.

We will not give any details concerning this derivation (we refer the interested reader to the work of H. Spohn [24]). We just claim that (formally at least) an appropriately rescaled Wigner transform of the displacement field q converges when $\lambda \rightarrow 0$ to a function $W(t, x, k)$ solution of the Boltzmann phonon equation

$$\partial_t W + \nabla_k \omega(k) \cdot \nabla_x W = C(W). \quad (3)$$

The function W depends on the time $t \geq 0$, the position $x \in \mathbb{R}^n$ and a wave vector k which lies in the Torus $\mathbb{T}^n = \mathbb{R}^n / \mathbb{Z}^n$. The function $\omega(k)$ is the dispersion relation of the lattice. It is determined by the harmonic part of the potential. For general potential given by (2), we have:

$$\omega(k) = (\omega_0^2 + \hat{\alpha}(k))^{1/2} \quad (4)$$

where $\widehat{\alpha}(k)$ is the Fourier transform of α , defined by

$$\widehat{\alpha}(k) = \sum_{j \in \mathbb{Z}^n} e^{-i2\pi k \cdot j} \alpha(j).$$

The operator C in the right hand side of (3) is an integral collision operator which depends on the anharmonic potential $V(q)$. Of course this operator C is crucial in determining the long time behavior of the solutions of this equation, so we will spend a bit of time discussing its properties in this introduction.

Note that while the relation between $W(t, x, k)$ and the microscopic variable q_i and p_i is rather complicated, the total energy of the system is given by

$$\begin{aligned} \int_{\mathbb{R}^n} \int_{\mathbb{T}^n} \omega(k) W(t, x, k) dk dx &= \frac{1}{2} \int |\widehat{p}(k)|^2 + \omega(k)^2 |\widehat{q}(k)|^2 dk \\ &= \sum_{i \in \mathbb{Z}^n} \frac{1}{2} p_i^2 + V_h(q). \end{aligned} \quad (5)$$

2.1 The FPU framework

As explained in the introduction, we now focus on the FPU chain model. For this model, we have $n = 1$ (we denote by \mathbb{T} the torus $\mathbb{T} = \mathbb{R}/\mathbb{Z}$) and the potential describes only nearest neighbors interactions. The harmonic potential is thus given by:

$$V_h(q) = \frac{1}{8} \sum_{i \in \mathbb{Z}} (q_{i+1} - q_i)^2,$$

and the anharmonic potential V is either cubic (FPU- α chain) or quartic (FPU- β chain):

$$V(q) = \sum_{i \in \mathbb{Z}^n} \gamma(q_{i+1} - q_i), \quad \gamma(q) = \frac{1}{3} q^3 \text{ or } \gamma(q) = \frac{1}{4} q^4.$$

The corresponding microscopic dynamics is given by

$$\begin{aligned} \frac{d}{dt} q_i(t) &= p_i(t) \\ \frac{d}{dt} p_i(t) &= \frac{1}{4} q_{i+1}(t) - \frac{1}{2} q_i(t) + \frac{1}{4} q_{i-1}(t) - \sqrt{\lambda} [\gamma'(q_i - q_{i-1}) - \gamma'(q_{i+1} - q_i)]. \end{aligned} \quad (6)$$

2.2 The dispersion relation

When V_h is given by

$$V_h(q) = \frac{1}{2}\omega_0^2 \sum_{i \in \mathbb{Z}} q_i^2 + \frac{1}{8} \sum_{i \in \mathbb{Z}} (q_{i+1} - q_i)^2, \quad (7)$$

equation (4) gives the following formula for the dispersion relation:

$$\omega(k)^2 = \omega_0^2 + \frac{1}{2} - \frac{1}{4} \left(e^{i2\pi k} + e^{-i2\pi k} \right)$$

and so

$$\omega(k) = \left(\omega_0^2 + \frac{1}{2}(1 - \cos(2\pi k)) \right)^{1/2}, \quad k \in \mathbb{T}. \quad (8)$$

For the FPU model, we have $\omega_0 = 0$, and so the dispersion relation is given by

$$\omega(k) = \sqrt{\frac{1}{2}(1 - \cos(2\pi k))} = |\sin(\pi k)|.$$

2.3 The interaction operator C

The operator C in the right hand side of (3) is determined by the non-harmonic perturbation of the potential V .

Cubic potentials: Three phonons operator When the anharmonic potential is cubic, that is

$$V = \frac{1}{3} \sum_{i \in \mathbb{Z}} q_i^3, \quad (9)$$

or

$$V = \frac{1}{3} \sum_{i \in \mathbb{Z}} (q_{i+1} - q_i)^3 \quad (10)$$

(the latter one corresponds to the FPU- α chain), the collision operator is given by

$$\begin{aligned} C(W) = & 4\pi \int \int F(k, k_1, k_2)^2 \\ & \times \left[2\delta(k + k_1 - k_2)\delta(\omega + \omega_1 - \omega_2)(W_1W_2 + WW_2 - WW_1) \right. \\ & \left. + \delta(k - k_1 - k_2)\delta(\omega - \omega_1 - \omega_2)(W_1W_2 - WW_1 - WW_2) \right] dk_1 dk_2 \end{aligned} \quad (11)$$




Figure 1: Three phonons interactions

where we used the notation $\omega_i = \omega(k_i)$ and $W_i = W(k_i)$.

The formula for the collision rate $F(k, k_1, k_2)$ can be found in [24]. In particular, when V is given by (9) (on-site potential) then

$$F(k, k_1, k_2)^2 = (8\omega\omega_1\omega_2)^{-1}$$

When V is the nearest neighbor interaction potential (10) and $\omega_0 = 0$ (that is for the FPU- α chain), the collision rate becomes

$$F(k, k_1, k_2)^2 = (8\omega\omega_1\omega_2)^{-1} |\exp(i2\pi k) - 1| |\exp(i2\pi k_1) - 1| |\exp(i2\pi k_2) - 1|^2.$$

Using the fact that

$$|\exp(i2\pi k) - 1|^2 = 4 \sin^2(\pi k),$$

we find

$$F(k, k_1, k_2)^2 = 8 \frac{\sin^2(\pi k) \sin^2(\pi k_1) \sin^2(\pi k_2)}{\omega\omega_1\omega_2}.$$

Going back to (11), we note that the first term can be interpreted as describing a wave vector k merging with a wave vector k_1 and leading to a new wave vector k_2 ($k + k_1 \rightarrow k_2$), while the second term describes the splitting of wave vector k into k_1 and k_2 ($k \rightarrow k_1 + k_2$). See Figure 1. These interactions conserve the energy ($\omega + \omega_1 = \omega_2$), but the momentum is conserved only modulo integers: the δ -function in the first term yields the constraint $k + k_1 = k_2 + n$, $n \in \mathbb{Z}$, $k, k_1, k_2 \in \mathbb{T}$ (one talks of normal process when $n = 0$, and *umklapp* process when $n \neq 0$).

This quadratic operator is reminiscent of the Boltzmann operator for the theory of dilute gas. There is however an essential difference: The kinetic energy $\frac{1}{2}v^2$ is replaced here by the dispersion relation $\omega(k)$. In order to further study this integral operator, it is thus essential to characterize the

set of (k, k_1, k_2) such that the δ -functions are not zero, that is:

$$\begin{cases} k + k_1 = k_2 \\ \omega(k) + \omega(k_1) = \omega(k_2) \end{cases}$$

or

$$\omega(k) + \omega(k_1) = \omega(k + k_1), \quad (k, k_1) \in \mathbb{T}, \quad (12)$$

This is much more delicate than for the usual Boltzmann operator and for general dispersion relation ω , it is not obvious that (12) has any solutions.

In our framework, that is when ω is given by (8) (nearest neighbor harmonic coupling) we actually can prove that

$$\omega(k) + \omega(k_1) - \omega(k + k_1) \geq \frac{\omega_0}{2}$$

so (12) has no solutions when $\omega_0 > 0$ and only the trivial solution $k_1 = 0$ when $\omega_0 = 0$.

It follows [24]:

Theorem 2.1. *When ω is given by (8) with $\omega_0 \geq 0$, then the three phonon collision operator (11) satisfies $C(W) = 0$ for all W .*

In particular, this implies that for the FPU- α chain, the collision operator vanishes, and the corresponding Boltzmann phonon equation reduces to pure transport. This suggests poor relaxation to equilibrium for the microscopic model, and it means that this kinetic approach is of no use in studying the long time behavior of the hamiltonian system. This is of course the reason why we focus in this paper on the FPU- β chain.

Remarks 2.2. *As noted in [24], equation (12) might have non trivial solutions for other dispersion relations (for instance $\omega(k) = \omega_0 + 2(1 - \cos(2\pi k))$), so this three phonon operator is of interest in other framework (different harmonic potential V_h).*

Quartic potentials: Four phonons operator. We now consider the quartic potential given by

$$V(q) = \frac{1}{4} \sum_{i \in \mathbb{Z}} q_i^4 \quad (13)$$




Figure 2: Four phonons interactions

or

$$V = \frac{1}{4} \sum_{i \in \mathbb{Z}} (q_{i+1} - q_i)^4. \quad (14)$$

The corresponding collision operator then reads

$$\begin{aligned} C(W) = 12\pi \sum_{\sigma_1, \sigma_2, \sigma_3 = \pm 1} \int \int \int F(k, k_1, k_2, k_3)^2 \\ \times \delta(k + \sigma_1 k_1 + \sigma_2 k_2 + \sigma_3 k_3) \delta(\omega + \sigma_1 \omega_1 + \sigma_2 \omega_2 + \sigma_3 \omega_3) \\ \times (W_1 W_2 W_3 + W(\sigma_1 W_2 W_3 + W_1 \sigma_2 W_3 + W_1 W_2 \sigma_3)) dk_1 dk_2 dk_3 \end{aligned} \quad (15)$$

with

$$F(k, k_1, k_2, k_3)^2 = (16\omega\omega_1\omega_2\omega_3)^{-1}$$

for on-site potential (13) and

$$F(k, k_1, k_2, k_3)^2 = \prod_{i=0}^3 \frac{2 \sin^2(\pi k_i)}{\omega(k_i)}. \quad (16)$$

for nearest neighbor coupling (14).

The term proportional to W is the loss term, while the gain term is $W_1 W_2 W_3$ (which is always positive). Again, we can interpret the different terms as pair collisions or merging/splitting of phonons (see Figure 2). In order to understand the collision rule, we note that for pair collisions $(k, k_1) \rightarrow (k_2, k_3)$ (which correspond to the terms such that $\sum_{j=1}^3 \sigma_j = -1$ in the integral), we need to solve

$$\omega(k) + \omega(k_1) = \omega(k_2) + \omega(k + k_1 - k_2) \quad (17)$$

while for three phonons mergers (or splitting) $(k, k_1, k_2) \rightarrow k_3$ we have

$$\omega(k) + \omega(k_1) + \omega(k_2) = \omega(k + k_1 + k_2). \quad (18)$$

In general, it is not possible to solve these equations explicitly, and it is not obvious that either of these equations should be satisfied on a set of positive measure. In fact, when ω is given by (8) (nearest neighbor couplings), it can be shown (see [25]) that (18) has no solution (so collision processes in which three phonons are merged into one, or one phonon splits into three are impossible). As a consequence, the only interactions that are allowed are pair collisions, which, in particular, preserve the total number of phonons. This preservation of the number of phonons, reminiscent of the preservation of the number of particles in gas dynamics, does not follow here from a fundamental physical principle, but is instead a mathematical artifact. This property is however stable under small perturbation of ω , and it also holds for the nonlinear wave equation for which $\omega(k) = |k|$ ($k \in \mathbb{R}^3$).

As a consequence, the operator C can be rewritten as

$$C(W) = 36\pi \int \int \int F(k, k_1, k_2, k_3)^2 \delta(k + k_1 - k_2 - k_3) \delta(\omega + \omega_1 - \omega_2 - \omega_3) [W_1 W_2 W_3 + W W_2 W_3 - W W_1 W_3 - W W_1 W_2] dk_1 dk_2 dk_3. \quad (19)$$

When ω is given by (8), we will see later on that (17) has non trivial solutions on a set of full measure, that is

$$\int_{\mathbb{T}} \int_{\mathbb{T}} \delta(\omega(k) + \omega(k_1) - \omega(k_2) - \omega(k + k_1 - k_2)) dk_1 dk_2 \neq 0.$$

In particular this operator C is non trivial.

3 FPU- β chain: The four phonon collision operator

In this section, we briefly summarize the properties of the four phonon collision operator (19) which arises in the modeling of the FPU- β chain.

3.1 Conserved quantities

All the collision operators C mentioned above conserve the energy. This can be expressed by the following condition:

$$\int_{\mathbb{T}} \omega(k) C(W)(k) dk = 0$$

for all functions W .

The four phonon collision operator (19), corresponding to the quartic potential, also satisfies

$$\int_{\mathbb{T}} C(W)(k) dk = 0$$

which can be interpreted as the conservation of the number of phonons $\int_{\mathbb{T}} W dk$. However, this quantity has no microscopic equivalent, and does not correspond to any physical principle. Rather it is a consequence of the symmetry of the operator, which follows from the fact that 3 phonon merger cannot take place ((18) has no solutions). In particular, this equality does not hold for the three phonon operator.

Note that the first moment k is preserved in the wave kinetic equation case (where $k \in \mathbb{R}^n$). However, this conservation is broken here by umklapp processes.

3.2 Entropy

The Boltzmann phonon operators satisfy an entropy inequality, similar to Boltzmann H-Theorem in gas dynamic. In particular, for the four phonon operator we can rewrite (19) as follows:

$$C(W) = 36\pi \int \int \int F(k)^2 \delta(k + k_1 - k_2 - k_3) \delta(\omega + \omega_1 - \omega_2 - \omega_3) \\ WW_1 W_2 W_3 [W^{-1} + W_1^{-1} - W_2^{-1} - W_3^{-1}] dk_1 dk_2 dk_3$$

and we then see that (assuming all integrals are well defined):

$$\int_{\mathbb{T}^1} W^{-1}(k) C(W)(k) dk \tag{20} \\ = 9\pi \int \int \int \int F(k)^2 \delta(k + k_1 - k_2 - k_3) \delta(\omega + \omega_1 - \omega_2 - \omega_3) \\ \cdot WW_1 W_2 W_3 [W^{-1} + W_1^{-1} - W_2^{-1} - W_3^{-1}]^2 dk_1 dk_2 dk_3 \\ \geq 0.$$

3.3 Stationary solutions

It is easy to check that the distributions

$$W_b(k) = \frac{1}{b\omega(k)}$$

for any $b > 0$ satisfy $C(W_b) = 0$ for all the operators C considered above. This fact is in accordance with equilibrium statistical mechanics (see [24]).

It is more delicate to check that these are the only solutions. In fact it is not always true.

For the four phonon collision operator (19), we can check that

$$W_{a,b}(k) = \frac{1}{a + b\omega(k)} \quad (21)$$

is an equilibrium for all $a, b > 0$.

Conversely, the entropy inequality (20) implies that if $C(W) = 0$ then $\psi(k) = W(k)^{-1}$ is a collision invariant, that is

$$\psi(k) + \psi(k_1) = \psi(k_2) + \psi(k_3)$$

for all k, k_1, k_2, k_3 such that

$$k + k_1 = k_2 + k_3, \quad \text{and} \quad \omega(k) + \omega(k_1) = \omega(k_2) + \omega(k_3).$$

An obvious candidate is $\psi(k) = a + b\omega(k)$. Under general conditions on ω , Spohn proved that these are indeed the only collision invariants in dimension $N \geq 2$ [23]. The same result is proved by Lukkarinen and Spohn [15] in our framework (dimension 1).

As a conclusion, (21) are the only solutions of $C(W) = 0$ for the four phonon collision operator (19). Note that the fact that we can take $a \neq 0$ is a consequence of the conservation of the number of phonons for the four phonon collision operator (which, as explained above, follows from the fact that equation (18) describing merging and splitting of phonons has no solutions).

3.4 The linearized operator

As mentioned in the introduction, we will be interested in the behavior of the solutions of the Boltzmann phonon equation in the neighborhood of a thermodynamical equilibrium. Given $\bar{W}(k) = \frac{T}{\omega(k)}$, we thus introduce the linearized operator

$$L(f) = \frac{1}{\bar{W}} DC(\bar{W})(\bar{W}f)$$

(where DC denotes the derivative of the operator C).

By differentiating the equation $C(W_{a,b}) = 0$ with respect to a and b , we get:

$$L(1) = 0 \quad \text{and} \quad L(\omega^{-1}) = 0,$$

which suggests (as will be proved later) that the kernel of L is two dimensional and spanned by 1 and ω^{-1} . In our framework, the later mode, ω^{-1} is singular (not integrable) for $k = 0$. Because of natural a priori bounds on the solutions of the Boltzmann Phonon equation, it will be easy to see that this mode is not present in the macroscopic limit. It will however play an important role in the derivation of a macroscopic model. Note that it comes from the derivation with respect to the spurious coefficient a .

Similarly, differentiating the conservation equations

$$\int \omega C(\bar{W} + t\bar{W}f) dk = 0 \text{ and } \int C(\bar{W} + t\bar{W}f) dk = 0$$

with respect to t , we deduce that

$$\int L(f) dk = 0, \text{ and } \int \omega^{-1}L(f) dk = 0.$$

The properties of L will be further investigated in Section 5. For now, we just state the following proposition without proof, since it is all we need to formally derive a macroscopic equation.

- Proposition 3.1.** *The operator $L : L^2(\mathbb{T}^1, V(k) dk) \longrightarrow L^2(\mathbb{T}^1, V(k)^{-1} dk)$ (where V is defined by (32)) is a bounded self-adjoint operator which satisfies*
- (i) $\ker(L) = \text{Span}\{1, \omega(k)^{-1}\}$
 - (ii) $R(L) = \{h \in L^2(\mathbb{T}^1, V(k)^{-1} dk); \int_{\mathbb{T}} h(k) dk = \int_{\mathbb{T}} \omega^{-1}(k)h(k) dk = 0\}$

We end this section by deriving the explicit formula for the operator L : A direct computation gives (when $W(k) = \frac{\bar{T}}{\omega(k)}$):

$$\begin{aligned} DC(W)(Wf) &= 36\pi \int \int \int F(k, k_1, k_2, k_3)^2 \delta(k + k_1 - k_2 - k_3) \delta(\omega + \omega_1 - \omega_2 - \omega_3) \\ &\quad \times WW_1W_2W_3 \left[f_3W_3^{-1} + f_2W_2^{-1} - f_1W_1^{-1} - fW^{-1} \right] dk_1 dk_2 dk_3 \\ &= 36\pi \bar{T}^3 \int \int \int \frac{F(k, k_1, k_2, k_3)^2}{\omega\omega_1\omega_2\omega_3} \delta(k + k_1 - k_2 - k_3) \delta(\omega + \omega_1 - \omega_2 - \omega_3) \\ &\quad \times \left[\omega_3f_3 + \omega_2f_2 - \omega_1f_1 - \omega f \right] dk_1 dk_2 dk_3 \end{aligned}$$

Using (16), we see that

$$\frac{F(k, k_1, k_2, k_3)^2}{\omega\omega_1\omega_2\omega_3} = 16$$

and we deduce:

$$L(f) = 576\pi\bar{T}^2\omega \int \int \int \delta(k + k_1 - k_2 - k_3)\delta(\omega + \omega_1 - \omega_2 - \omega_3) \\ \times \left[\omega_3 f_3 + \omega_2 f_2 - \omega_1 f_1 - \omega f \right] dk_1 dk_2 dk_3. \quad (22)$$

3.5 Formal asymptotic limit

We now have all the ingredient to perform the usual asymptotic analysis and attempt to derive (formally) a diffusion equation from the Boltzmann phonon equation (we will see however that it fails in our framework). The starting point is the following rescaled equation in the FPU- β chain framework detailed above:

$$\varepsilon^2 \partial_t W + \varepsilon \omega'(k) \partial_x W = C(W), \quad (23)$$

where C is the four phonon collision operator (19) with collision frequency given by (16), and we consider a solution which is a perturbation of a thermodynamical equilibrium:

$$W^\varepsilon(t, x, k) = \bar{W}(k)(1 + \varepsilon f^\varepsilon(t, x, k))$$

where $\bar{W} = \frac{\bar{T}}{\omega(k)}$ for some constant $\bar{T} > 0$.

We introduce the operators

$$Q(f, f) = \frac{1}{\bar{W}} D^2 C(\bar{W})(\bar{W}f, \bar{W}f),$$

and

$$R(f, f, f) = \frac{1}{\bar{W}} D^3 C(\bar{W})(\bar{W}f, \bar{W}f, \bar{W}f)$$

so that (we recall that C is a cubic operator):

$$\frac{1}{\bar{W}} C(W^\varepsilon) = \varepsilon L(f) + \varepsilon^2 \frac{1}{2} Q(f, f) + \varepsilon^3 \frac{1}{6} R(f, f, f)$$

where L is given by (22).

The function f^ε solves

$$\varepsilon^2 \partial_t f^\varepsilon + \varepsilon \omega'(k) \partial_x f^\varepsilon = L(f^\varepsilon) + \varepsilon \frac{1}{2} Q(f^\varepsilon, f^\varepsilon) + \varepsilon^2 \frac{1}{6} R(f^\varepsilon, f^\varepsilon, f^\varepsilon). \quad (24)$$

Taking the limit $\varepsilon \rightarrow 0$ in (24), we get

$$L(f^0) = 0$$

and so Proposition 3.1 (i) implies

$$f^0(t, x, k) = T(t, x) + S(t, x)\omega(k)^{-1}.$$

Since equation (24) preserves the L^1 norm, it is natural to assume that $f^0(t, x, k) \in L^1(\mathbb{R} \times \mathbb{T})$. We note however that $\omega(k) \sim |k|$ as $|k| \rightarrow 0$, and so we must have

$$S(t, x) = 0.$$

Next, integrating (24) with respect to k yields

$$\partial_t T^\varepsilon + \partial_x J^\varepsilon = 0$$

with

$$T^\varepsilon = \langle f^\varepsilon \rangle, \quad J^\varepsilon(t, x) = \frac{1}{\varepsilon} \langle \omega' f^\varepsilon \rangle$$

where we use the notation $\langle \cdot \rangle = \int_{\mathbb{T}} \cdot dk$.

We now need to compute $J = \lim_{\varepsilon \rightarrow 0} J^\varepsilon$. Recalling that L is a self adjoint operator, we write

$$\varepsilon^{-1} \langle \omega' f^\varepsilon \rangle = \langle L^{-1}(\omega') L(f^\varepsilon) \rangle$$

and using (24), we replace $L(f^\varepsilon)$ in the right hand side:

$$\varepsilon^{-1} \langle \omega' f^\varepsilon \rangle = \langle L^{-1}(\omega') \omega' \partial_x f^\varepsilon \rangle - \langle L^{-1}(\omega') Q(f^\varepsilon, f^\varepsilon) \rangle + \mathcal{O}(\varepsilon).$$

Formally, we thus get

$$\lim_{\varepsilon \rightarrow 0} \varepsilon^{-1} \langle \omega' f^\varepsilon \rangle = \langle L^{-1}(\omega') \omega' \rangle \partial_x T - \langle L^{-1}(\omega') Q(T, T) \rangle.$$

Finally, a direct computation gives

$$Q(f, f) = 576\pi \bar{T}^2 \omega \int \int \int \delta(k + k_1 - k_2 - k_3) \delta(\omega + \omega_1 - \omega_2 - \omega_3) \\ \left[2(\omega - \omega_3)[f_1 f_2 - f f_3] + (\omega + \omega_1)[f_2 f_3 - f f_1] \right] dk_1 dk_2 dk_3,$$

and it is readily seen that $Q(T, T) = 0$. We thus get the following relation

$$J = \langle L^{-1}(\omega') \omega' \rangle \partial_x T$$

which is Fourier's law with diffusion coefficient

$$\kappa = -\langle L^{-1}(\omega') \omega' \rangle > 0.$$

We conclude this section with the following remarks:

1. The non linear term $Q(T, T) = 0$ does not contribute to the limiting equation. In the next section, we will drop this term altogether.
2. The fact that $S = 0$ will need to be addressed very carefully in the rigorous proof. In particular, we will see that while we do indeed have $f^0 = T$, the term S plays a significant role in the rigorous derivation of the diffusion equation (see next section).
3. Perhaps the most important remark is that one need to check that κ is well defined. In fact, it can be proved that the integrand in the definition of the diffusion coefficient behaves like $|k|^{-5/3}$ for small k . It follows that

$$\kappa = +\infty$$

so the limit presented above does not give any equation for the evolution of T . Such a phenomenon is not uncommon, and based on previous work (see [17]), we expect that by taking a different time scale in (24) we can derive an anomalous diffusion equation for the evolution of the temperature T . This is of course the goal of this paper as explained in the next section.

4 Main result

In view of the formal asymptotic limit detailed in the previous section, we now consider the following linear equation:

$$\varepsilon^\alpha \partial_t f^\varepsilon + \varepsilon \omega'(k) \partial_x f^\varepsilon = \bar{T}^2 L(f^\varepsilon), \quad x \in \mathbb{R}, k \in \mathbb{T} \quad (25)$$

where

$$\omega(k) = |\sin(\pi k)|$$

and L is defined by

$$L(f) = \omega \int \int \int \delta(k + k_1 - k_2 - k_3) \delta(\omega + \omega_1 - \omega_2 - \omega_3) \left[\omega_3 f_3 + \omega_2 f_2 - \omega_1 f_1 - \omega f \right] dk_1 dk_2 dk_3. \quad (26)$$

Note also that we have made L independent of the equilibrium temperature \bar{T} and set all other constant in L equal to 1 for the sake of clarity.

The existence of a solution to this equation is fairly classical. We recall it for the sake of completeness in Proposition 5.10.

Our main result is then the following:

Theorem 4.1 (Fractional diffusion limit for the linearised equation). *Take $\alpha = \frac{8}{5}$ and let f^ε be a solution of equation (25) with initial data $f_0 \in L^2(\mathbb{R} \times \mathbb{T})$. Then*

$$f^\varepsilon(t, x, k) \rightharpoonup T(t, x) \quad L^\infty((0, \infty); L^2(\mathbb{R} \times \mathbb{T}))\text{-weak*}$$

where T solves the fractional diffusion equation

$$\partial_t T + \frac{\kappa}{\bar{T}^{6/5}} (-\Delta_x)^{4/5} T = 0 \quad \text{in } (0, \infty) \times \mathbb{R} \quad (27)$$

with initial condition

$$T(0, x) = T_0(x) := \int_0^1 f_0(x, k) dk. \quad (28)$$

The diffusion coefficient $\kappa \in (0, \infty)$ is given by

$$\kappa = \kappa_1 - \frac{\kappa_2^2}{\kappa_3} \in (0, \infty)$$

where $\kappa_1, \kappa_2, \kappa_3$ are defined in Proposition 6.4.

First, we note that it is enough to consider the case

$$\bar{T} = 1$$

since we can recover the general case by a simple rescaling $t \mapsto \bar{T}^2 t, x \mapsto \bar{T}^2 x$.

The main difficulty here, compared with previous work devoted to fractional diffusion limit of kinetic equations, is the fact that the kernel of L is spanned by 1 and $\omega(k)^{-1}$. This last mode should not appear in the limit since it is not square integrable, but it will nevertheless play an important role.

In fact, we will prove that f^ε can be expanded as follows:

$$f^\varepsilon(t, x, k) = T^\varepsilon(t, x) + \varepsilon^{\frac{3}{5}} S^\varepsilon(t, x) \omega(k)^{-1} + \varepsilon^{\frac{4}{5}} h^\varepsilon(t, x, k)$$

where T^ε is bounded in $L^\infty(0, \infty; L^2(\mathbb{R}))$, h^ε is bounded in $L_V^2(\mathbb{T} \times \mathbb{R})$ and S^ε converges in some weak sense to a non trivial function. More precisely we will prove in Section 7:

Proposition 4.2. *The function $S^\varepsilon(t, x)$ converges in distribution sense to*

$$S(t, x) = -\frac{\kappa_2}{\kappa_3} (-\Delta)^{3/10} T(t, x).$$

In particular, as mentioned above, this means that the mode $\omega(k)^{-1}$ vanishes in the limit and the macroscopic behavior of the phonon distribution is completely described by $T = \lim_{\varepsilon \rightarrow 0} T^\varepsilon$. However, projecting equation (25) onto the constant mode of the kernel of L , we will find the following equation of the evolution of T :

$$\partial_t T + \kappa_1(-\Delta)^{4/5}T + \kappa_2(-\Delta)^{1/2}S = 0. \quad (29)$$

We see that $S = \lim_{\varepsilon \rightarrow 0} S^\varepsilon$ plays a role in the evolution of T . To understand this, we note (anticipating a bit on the result of the next section) that the reason we are observing anomalous diffusion phenomena here (as opposed to standard diffusion as described in the previous section), is the fact that phonons with frequency k close to zero encounter very few collisions (degenerate collision frequency). And the term $\varepsilon^{\frac{3}{5}}S^\varepsilon(t, x)\omega(k)^{-1}$, while small, is heavily concentrated around $k = 0$ (non integrable singularity at $k = 0$). The competition between the smallness and the singularity gives rise to a term of order 1 in the equation.

In order to describe the evolution of T , we now need to obtain an equation for S . By projecting equation (25) onto the $\omega(k)^{-1}$ mode of the kernel of L , we will prove that:

$$\kappa_2(-\Delta)^{1/2}T + \kappa_3(-\Delta)^{1/5}S = 0. \quad (30)$$

We note that there is no $\partial_t S$ in (30) (unlike the corresponding equation for T). The reason is that due to the singularity of $\omega(k)^{-1}$ for $k = 0$, the quantity S diffuses faster than T (so we would have to take a smaller α in (25) in order to observe the diffusion of S). At our time scale (given by $\alpha = \frac{8}{5}$), S has thus already reached equilibrium, and can be expressed (in view of (30)) as

$$S = -\frac{\kappa_2}{\kappa_3}(-\Delta)^{3/10}T.$$

Inserting this expression into (29), we find

$$\partial_t T + \kappa(-\Delta)^{4/5}T = 0$$

where $\kappa = \kappa_1 - \frac{\kappa_2^2}{\kappa_3}$. Of course, we will show that $\kappa > 0$ (once the explicit expressions for the κ_i are given, it will be a very simple consequence of Cauchy-Schwarz inequality - see Lemma 6.6). It is interesting to note that the effect of the mode ω^{-1} on the macroscopic equation is to reduce the diffusion coefficient (and thus to slow down the diffusion). This can be understood by noting that the fact that the kernel of L does not contain

only the natural constant mode, is due to the lack of merging $k+k_1+k_2 \rightarrow k_3$ and splitting $k \rightarrow k_1+k_2+k_3$ interactions for phonons in the non linear collision operator C (fewer interactions \Rightarrow slower relaxation).

5 Properties of the operator L

The asymptotic behavior of the solution of (25) depends very strongly on the properties of the operator L . This operator is studied in great detail in [15], and we will recall their main results in this section.

The operator L can be written as

$$L(f) = \int K(k, k') f(k') dk' - V(k) f(k)$$

where

$$\begin{aligned} K(k, k') = \omega(k)\omega(k') \int_{\mathbb{T}} & 2 \delta(\omega(k) + \omega(k_1) - \omega(k') - \omega(k + k_1 - k')) \\ & - \delta(\omega(k) + \omega(k') - \omega(k_1) - \omega(k + k' - k_1)) dk_1 \end{aligned} \quad (31)$$

and

$$V(k) = \omega(k)^2 \int_{\mathbb{T}} \delta(\omega(k) + \omega(k_1) - \omega(k') - \omega(k + k_1 - k')) dk_1 dk'. \quad (32)$$

The fact that $\int_{\mathbb{T}} L(f) dk = 0$ for all f implies

$$V(k) = \int_{\mathbb{T}} K(k', k) dk'$$

(this equality can be checked also from the formula for K and V , but it is much easier this way) and a short computation shows that

$$K(k, k') = K(k', k).$$

In particular, L is a self adjoint operator in $L^2(\mathbb{T})$. It is also positive since we have

$$\begin{aligned} - \int_{\mathbb{T}} L(f) f dk &= \frac{1}{4} \int \int \int \int \delta(k + k_1 - k_2 - k_3) \delta(\omega + \omega_1 - \omega_2 - \omega_3) \\ & [\omega_3 f_3 + \omega_2 f_2 - \omega_1 f_1 - \omega f]^2 dk dk_1 dk_2 dk_3 \quad (33) \\ &\geq 0 \end{aligned}$$

for all f . One of our goals will be to improve this inequality and show that L has a spectral gap property in the appropriate functional spaces. For that, we will need to show that the integral operator

$$K(f) = \int K(k, k')f(k') dk' \quad (34)$$

is a compact operator (in an appropriate functional spaces)

The first step, in view of (31) is to study the solution set of the equation of conservation of energy:

$$\omega(k) + \omega(k_1) = \omega(k') + \omega(k + k_1 - k'). \quad (35)$$

We recall the following result:

Proposition 5.1 ([15]). *The equation (35) has the trivial solutions $k' = k$ and $k' = k_1$, and the (non trivial) solution*

$$k_1 = h(k, k')$$

where

$$h(k, k') = \frac{k' - k}{2} + 2 \arcsin \left(\tan \frac{|k' - k|}{4} \cos \frac{k + k'}{4} \right)$$

(and there are no other solutions of (35)).

With this proposition in hand, one can now compute the kernel $K(k, k')$ and the multiplicative function $V(k)$. We recall here the main result of [15]. The first one states that the function $V(k)$ is degenerate for $k \rightarrow 0$ (note that W in [15] corresponds to our V):

Proposition 5.2 ([15, Lemma 4.1]). *The function $V : \mathbb{R} \rightarrow \mathbb{R}_+$ is symmetric ($V(1 - k) = V(k)$), continuous and satisfies*

$$c_1 |\sin(\pi k)|^{5/3} \leq V(k) \leq c_2 |\sin(\pi k)|^{5/3} \quad (36)$$

for all $k \in \mathbb{R}$, for some $c_1, c_2 > 0$. Moreover,

$$\lim_{k \rightarrow 0} \left(|\sin \pi k|^{-5/3} V(k) \right) = v_0 > 0.$$

Because of the degeneracy of V for $k = 0$, we do not expect the operator L to have a spectral gap in L^2 . We thus introduce the operator

$$L_0(f) := V^{-1/2} L(V^{-1/2} f)$$

We note that this operator has the form

$$L_0(f) = K_0(f) - f$$

with

$$K_0(f) = V^{-1/2}K(V^{-1/2}f).$$

To prove that L_0 has good properties in $L^2(\mathbb{T})$, we need to study the operator K_0 . Again, it is proved in [15] that $K_0 : L^2(\mathbb{T}^1) \rightarrow L^2(\mathbb{T}^1)$ is a compact, self-adjoint operator, which implies that $K : L^2(\mathbb{T}^1, V dk) \rightarrow L^2(\mathbb{T}^1, V^{-1}dk)$ is a compact, self-adjoint operator.

To be more precise, in [15], the kernel K is first written as

$$K(k, k') = 2\omega(k)K_2(k, k')\omega(k') - \omega(k)K_1(k, k')\omega(k')$$

where

$$K_1(k, k') := 4 \frac{\mathbb{1}(F_-(k, k') > 0)}{\sqrt{F_-(k, k')}} \quad \text{and} \quad K_2(k, k') := \frac{2}{\sqrt{F_+(k, k')}} \quad (37)$$

for $k, k' \in [0, 1]$ and

$$F_{\pm}(k, k') = (\cos(\pi k) + \cos(\pi k'))^2 \pm 4 \sin(\pi k) \sin(\pi k').$$

and the main result of [15] is the following:

Proposition 5.3 ([15, Propositions 4.3 and 4.4.]). *Let $\psi : [0, 1] \rightarrow \mathbb{R}$ be given, and assume that there are $C, p > 0$ such that*

$$|\psi(k)| \leq C (\sin \pi k)^p$$

for all $k \in [0, 1]$. Then the kernels

$$\psi(k)^* K_2(k, k') \psi(k') \quad \text{and} \quad \psi(k)^* K_1(k, k') \psi(k')$$

define compact, self-adjoint integral operators in $L^2(\mathbb{T})$.

We immediately conclude:

Corollary 5.4. *The kernel*

$$K_0(k, k') = V^{-1/2}(k)\omega(k) (2K_2(k, k') - K_1(k, k')) \omega(k')V^{-1/2}(k') \quad (38)$$

defines a compact self-adjoint operator in $L^2((0, 1))$. As a consequence, the kernel

$$K(k, k') = V^{1/2}(k)K_0(k, k')V^{1/2}(k')$$

defines a compact self-adjoint operator from $L^2(\mathbb{T}^1, V(k) dk)$ onto $L^2(\mathbb{T}^1, V(k)^{-1} dk)$. In particular,

$$\int_{\mathbb{T}} |K(f)(k)|^2 V(k)^{-1} dk \leq C \int_{\mathbb{T}} |f(k)|^2 V(k) dk. \quad (39)$$

Proof. Indeed, by Proposition 5.2 we have that

$$V^{-1/2}(k)\omega(k) \leq c_2 (\sin \pi k)^{1/6}$$

and the claim follows from Proposition 5.3. \square

Furthermore, we note that we have not used the full potential of Proposition (5.3). We can thus improve (39) as follows:

Corollary 5.5. *The kernel*

$$\widetilde{K}_0(k, k') := (\sin(\pi k))^{-1/6+\eta} K_0(x, k') (\sin(\pi k'))^{-1/6+\eta} \quad \eta > 0$$

defines a compact self-adjoint operator in $L^2((0, 1))$. In particular, for all $\eta > 0$, there exists $C(\eta)$ such that

$$\int_{\mathbb{T}} |K(f)(k)|^2 (\sin(\pi k))^{-\frac{1}{3}+\eta} V(k)^{-1} dk \leq C \int_{\mathbb{T}} |f(k)|^2 (\sin(\pi k))^{\frac{1}{3}-\eta} V(k) dk \quad (40)$$

Proof. Using Proposition 5.2 we have that

$$\begin{aligned} V^{-1/2}(k)\omega(k) (\sin \pi k)^{-1/6+\eta} &\leq c_2 (\sin \pi k)^{1/6} (\sin \pi k)^{-1/6+\eta} \\ &= c_2 (\sin \pi k)^\eta \end{aligned}$$

the claim follows from Proposition 5.3. \square

We have thus showed that $L : L^2(\mathbb{T}^1, V(k) dk) \longrightarrow L^2(\mathbb{T}^1, V(k)^{-1} dk)$ was a bounded operator. Next, we characterize the kernel of L : First, we note that given $f \in L^2(\mathbb{T}^1, V(k) dk)$, inequality (33) implies that if $L(f) = 0$ then

$$\begin{aligned} &\int \int \int \int \delta(k + k_1 - k_2 - k_3) \delta(\omega + \omega_1 - \omega_2 - \omega_3) \\ &\quad \times [\omega_3 f_3 + \omega_2 f_2 - \omega_1 f_1 - \omega f]^2 dk dk_1 dk_2 dk_3 = 0. \end{aligned}$$

So f must satisfy

$$\omega(k)f(k) + \omega(k_1)f(k_1) = \omega(k_2)f(k_2) + \omega(k + k_1 - k_2)f(k + k_1 - k_2)$$

whenever

$$\omega(k) + \omega(k_1) = \omega(k_2) + \omega(k + k_1 - k_2).$$

We also say that $\omega(k)f(k)$ must be a collision invariant. Such invariants have been characterized in [15]:

Theorem 5.6 ([15]). *A function $\psi \in L^1(\mathbb{T})$ is a collisional invariant if and only if there exists c_1 and c_2 such that*

$$\psi(k) = c_1 + c_2\omega(k).$$

As a consequence, we deduce:

Corollary 5.7. *The kernel of L is the two dimensional subspace of $L^2(\mathbb{T}^1, V(k) dk)$ spanned by the functions 1 and $\omega(k)^{-1}$ (note that both of those functions belongs to $L^2(\mathbb{T}^1, V(k) dk)$ thanks to (36))*

Finally, the compactness of K and inequality (33) implies

Lemma 5.8. *There exists $c_0 > 0$ such that*

$$- \int_{\mathbb{T}^1} L(f)f dk \geq c_0 \int V(k)|f - \Pi(f)|^2 dk$$

for all $f \in L^2(\mathbb{T}^1, V(k) dk)$, where $\Pi(f)$ denotes the orthogonal projection of f onto $\ker(L)$.

To summarize, we have thus showed:

Proposition 5.9. *The operator $L : L^2(\mathbb{T}^1, V(k) dk) \longrightarrow L^2(\mathbb{T}^1, V(k)^{-1} dk)$ is bounded and satisfies:*

1. *The kernel of L has dimension 2 and is spanned by 1 and $\frac{1}{\omega(k)}$.*
2. *For all $f \in L^2(\mathbb{T}^1, V(k) dk)$, we have*

$$\int_{\mathbb{T}^1} L(f) dk = 0 \quad \text{and} \quad \int_{\mathbb{T}^1} \frac{1}{\omega(k)} L(f) dk = 0. \quad (41)$$

3. *There exists $c_0 > 0$ such that*

$$- \int_{\mathbb{T}^1} L(f)f dk \geq c_0 \int V(k)|f - \Pi(f)|^2 dk$$

for all $f \in L^2(\mathbb{T}^1, V(k) dk)$, where $\Pi(f)$ denotes the orthogonal projection of f onto $\ker(L)$.

Note that the projection of f onto $\ker(L)$ can be written as

$$\Pi(f) = T + S [\langle V \rangle \omega(k)^{-1} - \langle V \omega^{-1} \rangle]$$

with

$$T = \frac{1}{\langle V \rangle} \int V(k) f(k) dk \text{ and } S = \frac{1}{m_0} \int \left[\langle V \rangle \frac{V(k)}{\omega(k)} - \langle V \omega^{-1} \rangle V(k) \right] f(k) dk$$

where $m_0 = \langle V \rangle^2 \langle V \omega^{-2} \rangle - \langle V \omega^{-1} \rangle^2 \langle V \rangle$ is a normalization constant. The operator Π is a continuous operator in $L^2(V(k) dk)$.

We finish this section commenting on the existence of solutions for the equation for the sake of completeness:

Proposition 5.10 (Cauchy Problem). *There exists a unique solution in $L^\infty((0, \infty); L^2(\mathbb{R} \times \mathbb{T}))$ for equation (25) with initial data $f_0 \in L^2(\mathbb{R} \times \mathbb{T})$.*

Proof. A traditional method for solving the Cauchy problem for this type of equations uses an iterative scheme based on the mild formulation:

$$f(t, x, k) = f_0(x - \omega'(k)t, k) + \int_0^t Lf(x - (t-s)\omega'(k), s) ds$$

together with the estimate

$$\|L(f)\|_{L^2(\mathbb{R} \times \mathbb{T})} \leq C \|f\|_{L^2(\mathbb{R} \times \mathbb{T})}.$$

This last estimate is consequence of (39) and the boundedness of the function V . We refer to [2] and [18] for further details on this method. \square

6 Proof of Theorem 4.1

6.1 A priori estimates

As a first step in the proof of Theorem 4.1, we establish some a priori estimates. The coercivity property of L (Lemma 5.8) gives the following proposition:

Proposition 6.1. *Assume that $f_0 \in L^2(\mathbb{R} \times \mathbb{T})$. Then, the function $f^\varepsilon(t, x, k)$, solution of (25) satisfies*

$$\|f^\varepsilon(t)\|_{L^2(\mathbb{R} \times \mathbb{T})} \leq \|f_0\|_{L^2(\mathbb{R} \times \mathbb{T})} \quad \text{for all } t \geq 0. \quad (42)$$

Furthermore, f^ε can be expanded as follows:

$$f^\varepsilon = \Pi(f^\varepsilon) + \varepsilon^{4/5} h^\varepsilon, \quad (43)$$

where

$$\|h^\varepsilon\|_{L_V^2((0,\infty)\times\mathbb{R}\times\mathbb{T})} \leq C\|f_0\|_{L^2(\mathbb{R}\times\mathbb{T})} \quad (44)$$

and $\Pi(f^\varepsilon)$ is the projection of f^ε onto $\ker(L)$, given by

$$\Pi(f^\varepsilon)(t, x, k) = \tilde{T}^\varepsilon(t, x) + \tilde{S}^\varepsilon(t, x) [\langle V \rangle \omega(k)^{-1} - \langle V \omega^{-1} \rangle]$$

with

$$\begin{aligned} \tilde{T}^\varepsilon(t, x) &= \frac{1}{\langle V \rangle} \int V(k) f^\varepsilon(t, x, k) dk, \\ \tilde{S}^\varepsilon(t, x) &= \frac{1}{m_0} \int \left[\langle V \rangle \frac{V(k)}{\omega(k)} - \langle V \omega^{-1} \rangle V(k) \right] f^\varepsilon(t, x, k) dk \end{aligned} \quad (45)$$

where $\tilde{T}^\varepsilon, \tilde{S}^\varepsilon$ are bounded in $L^\infty((0, \infty); L^2(\mathbb{R}))$.

Proof. Multiplying (25) by f^ε and integrating with respect to x and k , we get

$$\frac{1}{2} \frac{d}{dt} \|f^\varepsilon(t)\|_{L^2(\mathbb{R}\times\mathbb{T}^1)}^2 - \frac{1}{\varepsilon^\alpha} \int_{\mathbb{R}} \int_{\mathbb{T}^1} L(f^\varepsilon) f^\varepsilon dk dx = 0.$$

Integrating with respect to t and using (Lemma 5.8), we deduce

$$\frac{1}{2} \|f^\varepsilon(t)\|_{L^2(\mathbb{R}\times\mathbb{T}^1)}^2 + \frac{c_0}{\varepsilon^\alpha} \int_0^t \int_{\mathbb{R}} \int_{\mathbb{T}^1} V(k) |f^\varepsilon - \Pi(f^\varepsilon)|^2 dk dx ds \leq \frac{1}{2} \|f_0^\varepsilon\|_{L^2(\mathbb{R}\times\mathbb{T}^1)}^2.$$

which implies the proposition. The fact that $\tilde{T}^\varepsilon, \tilde{S}^\varepsilon \in L^\infty((0, \infty); L^2(\mathbb{R}))$ is a direct consequence of this estimate and Cauchy-Schwartz. \square

Because the singular terms in $\Pi(f^\varepsilon)$ (those involving $\omega(k)^{-1}$) play a particular role in the sequel, we will prefer to write $\Pi(f^\varepsilon)$ as follows:

$$\Pi f^\varepsilon = T^\varepsilon + \frac{\langle V \rangle}{\omega} \tilde{S}^\varepsilon(x, t)$$

with

$$T^\varepsilon(t, x) = \tilde{T}^\varepsilon(t, x) - \tilde{S}^\varepsilon(t, x) \langle V \omega^{-1} \rangle$$

Finally, we set

$$S^\varepsilon(t, x) = \varepsilon^{-3/5} \langle V \rangle \tilde{S}^\varepsilon(t, x), \quad (46)$$

leading to the following expansion of f^ε :

$$f^\varepsilon(t, x, k) = T^\varepsilon(t, x) + \varepsilon^{\frac{3}{5}} S^\varepsilon(t, x) \omega(k)^{-1} + \varepsilon^{\frac{4}{5}} h^\varepsilon(t, x, k). \quad (47)$$

Note that while T^ε and h^ε are clearly bounded (in appropriate functional spaces) in view of Proposition 6.1, the scaling of S^ε may seem arbitrary at this point. However, we will see later on that S^ε defined as in (46) indeed converges to a non trivial function (in some weak sense).

6.2 Laplace Fourier Transform

As in [17], the main tool in deriving the macroscopic equation for T is the use of the Laplace-Fourier transform. More precisely, we define

$$\widehat{f^\varepsilon}(p, \xi, k) = \int_{\mathbb{R}} \int_0^\infty e^{-pt} e^{-i\xi x} f^\varepsilon(t, x, k) dt dx.$$

We also denote by $\widehat{f_0}(\xi, k)$ the Fourier transform of $f_0(x, k)$.

Remarks 6.2. *We recall that the Fourier transform preserves the $L^2(\mathbb{R})$ norm (Parseval's theorem). It is also easy to see that the Laplace transform of an L^1 function is in L^∞ . However our functions are not L^1 with respect to t . Instead, we will make use of the simple fact that for a given function $g(t)$, its Laplace transform $\widehat{g}(p)$ satisfies*

$$|\widehat{g}(p)| \leq \frac{1}{p} \|g\|_{L^\infty(0, \infty)} \quad \text{and} \quad |\widehat{g}(p)| \leq \frac{1}{\sqrt{2p}} \|g\|_{L^2(0, \infty)} \quad (48)$$

for all $p > 0$.

Taking the Laplace Fourier transform of the equation, we obtain:

$$\varepsilon^\alpha p \widehat{f^\varepsilon} - \varepsilon^\alpha \widehat{f_0} + i\varepsilon \omega'(k) \xi \widehat{f^\varepsilon} = K(\widehat{f^\varepsilon}) - V \widehat{f^\varepsilon}$$

which easily yields

$$\widehat{f^\varepsilon}(p, \xi, k) = \frac{\varepsilon^\alpha}{\varepsilon^\alpha p + V(k) + i\varepsilon \omega'(k) \xi} \widehat{f_0} + \frac{1}{\varepsilon^\alpha p + V(k) + i\varepsilon \omega'(k) \xi} K(\widehat{f^\varepsilon}). \quad (49)$$

We recall that $L(f) = K(f) - Vf$ with $K(f) = \int K(k, k') f(k') dk'$. The fact that $\int L(f) dk = 0$ and $\int \frac{1}{\omega(k)} L(f) dk = 0$ for all f implies

$$V(k) = \int K(k', k) dk', \quad \frac{V(k)}{\omega(k)} = \int K(k', k) \frac{1}{\omega(k')} dk'$$

Multiplying (49) by $K(k', k)$ and integrating with respect to k and k' , we get

$$\begin{aligned}
\int_{\mathbb{T}} K(\widehat{f^\varepsilon})(k') dk' &= \int_{\mathbb{T}} \int_{\mathbb{T}} \frac{\varepsilon^\alpha K(k', k)}{\varepsilon^\alpha p + V(k) + i\varepsilon\omega'(k)\xi} \widehat{f_0}(\xi, k) dk dk' \\
&\quad + \int_{\mathbb{T}} \int_{\mathbb{T}} \frac{K(k', k)}{\varepsilon^\alpha p + V(k) + i\varepsilon\omega'(k)\xi} K(\widehat{f^\varepsilon})(k) dk dk' \\
&= \int_{\mathbb{T}} \frac{\varepsilon^\alpha V(k)}{\varepsilon^\alpha p + V(k) + i\varepsilon\omega'(k)\xi} \widehat{f_0}(\xi, k) dk \\
&\quad + \int_{\mathbb{T}} \frac{V(k)}{\varepsilon^\alpha p + V(k) + i\varepsilon\omega'(k)\xi} K(\widehat{f^\varepsilon})(k) dk.
\end{aligned}$$

We deduce

$$\begin{aligned}
0 &= \int_{\mathbb{T}} \frac{V(k)}{\varepsilon^\alpha p + V(k) + i\varepsilon\omega'(k)\xi} \widehat{f_0}(\xi, k) dk \\
&\quad + \varepsilon^{-\alpha} \int_{\mathbb{T}} \left(\frac{V(k)}{\varepsilon^\alpha p + V(k) + i\varepsilon\omega'(k)\xi} - 1 \right) K(\widehat{f^\varepsilon})(k) dk. \tag{50}
\end{aligned}$$

Similarly, multiplying (49) by $K(k', k) \frac{\varepsilon^{\frac{3}{5}}}{\omega(k')}$, and we get:

$$\begin{aligned}
0 &= \varepsilon^{\frac{3}{5}} \int_{\mathbb{T}} \frac{V(k)}{\varepsilon^\alpha p + V(k) + i\varepsilon\omega'(k)\xi} \frac{\widehat{f_0}(\xi, k)}{\omega(k)} dk \\
&\quad + \varepsilon^{-\alpha} \varepsilon^{\frac{3}{5}} \int_{\mathbb{T}} \left(\frac{V(k)}{\varepsilon^\alpha p + V(k) + i\varepsilon\omega'(k)\xi} - 1 \right) \frac{K(\widehat{f^\varepsilon})(k)}{\omega(k)} dk. \tag{51}
\end{aligned}$$

Next, we write

$$K(\widehat{f^\varepsilon}) = K(\Pi(\widehat{f^\varepsilon})) + K(\widehat{f^\varepsilon} - \Pi(\widehat{f^\varepsilon})) = V\Pi(\widehat{f^\varepsilon}) + K(\widehat{f^\varepsilon} - \Pi(\widehat{f^\varepsilon}))$$

where we rewrite

$$\Pi(\widehat{f^\varepsilon}) = \widehat{T}^\varepsilon + \varepsilon^{3/5} \frac{1}{\omega(k)} \widehat{S}^\varepsilon.$$

We can thus rewrite (50) as follows:

$$\mathcal{F}_1^\varepsilon(\widehat{f^0}) + a_1^\varepsilon(p, \xi) \widehat{T}^\varepsilon(p, \xi) + a_2^\varepsilon(p, \xi) \widehat{S}^\varepsilon(p, \xi) + R_1^\varepsilon(p, \xi) = 0 \tag{52}$$

and (51) as follows:

$$\mathcal{F}_2^\varepsilon(\widehat{f^0}) + a_2^\varepsilon(p, \xi) \widehat{T}^\varepsilon(p, \xi) + a_3^\varepsilon(p, \xi) \widehat{S}^\varepsilon(p, \xi) + R_2^\varepsilon(p, \xi) = 0, \tag{53}$$

where for $\alpha = 8/5$, we have:

$$\begin{aligned}\mathcal{F}_1^\varepsilon(\widehat{f}^0) &= \int_{\mathbb{T}} \frac{V(k)}{\varepsilon^{\frac{8}{5}}p + V(k) + i\varepsilon\omega'(k)\xi} \widehat{f}_0(\xi, k) dk \\ \mathcal{F}_2^\varepsilon(\widehat{f}^0) &= \varepsilon^{\frac{3}{5}} \int_{\mathbb{T}} \frac{V(k)}{\varepsilon^{\frac{8}{5}}p + V(k) + i\varepsilon\omega'(k)\xi} \frac{\widehat{f}_0(\xi, k)}{\omega(k)} dk,\end{aligned}$$

$$\begin{aligned}a_1^\varepsilon(p, \xi) &:= \varepsilon^{-\frac{8}{5}} \int_{\mathbb{T}} \left(\frac{V(k)}{\varepsilon^{\frac{8}{5}}p + V(k) + i\varepsilon\omega'(k)\xi} - 1 \right) V(k) dk \\ a_2^\varepsilon(p, \xi) &:= \varepsilon^{-\frac{8}{5}} \int_{\mathbb{T}} \left(\frac{V(k)}{\varepsilon^{\frac{8}{5}}p + V(k) + i\varepsilon\omega'(k)\xi} - 1 \right) \frac{\varepsilon^{\frac{3}{5}}V(k)}{\omega(k)} dk \\ &= \varepsilon^{-1} \int_{\mathbb{T}} \left(\frac{V(k)}{\varepsilon^{\frac{8}{5}}p + V(k) + i\varepsilon\omega'(k)\xi} - 1 \right) \frac{V(k)}{\omega(k)} dk \\ a_3^\varepsilon(p, \xi) &:= \varepsilon^{-1} \int_{\mathbb{T}} \left(\frac{V(k)}{\varepsilon^{\frac{8}{5}}p + V(k) + i\varepsilon\omega'(k)\xi} - 1 \right) \frac{\varepsilon^{\frac{3}{5}}V(k)}{\omega(k)^2} dk\end{aligned}$$

and

$$\begin{aligned}R_1^\varepsilon(\xi, p) &:= \varepsilon^{-\frac{8}{5}} \int_{\mathbb{T}} \left(\frac{V(k)}{\varepsilon^{\frac{8}{5}}p + V(k) + i\varepsilon\omega'(k)\xi} - 1 \right) K(\widehat{f}^\varepsilon - \Pi(\widehat{f}^\varepsilon))(k) dk \\ R_2^\varepsilon(\xi, p) &:= \varepsilon^{-1} \int_{\mathbb{T}} \left(\frac{V(k)}{\varepsilon^{\frac{8}{5}}p + V(k) + i\varepsilon\omega'(k)\xi} - 1 \right) \frac{1}{\omega(k)} K(\widehat{f}^\varepsilon - \Pi(\widehat{f}^\varepsilon))(k) dk\end{aligned}$$

In order to prove the main theorem, we now need to pass to the limit in (52) and (53). The following three propositions, which are proved in the next section, give the necessary results for that.

First, we have the following limits for the terms involving the initial data:

Proposition 6.3. *The following limits hold for all $P \geq 0$:*

$$\begin{aligned}\mathcal{F}_1^\varepsilon(\widehat{f}^0)(\xi, p) &\longrightarrow \int_{\mathbb{T}} \widehat{f}^0(\xi, k) dk = \widehat{T}_0(\xi) && \text{in } L^2((0, P) \times \mathbb{R}) \\ \mathcal{F}_2^\varepsilon(\widehat{f}^0)(\xi, p) &\longrightarrow 0 && \text{in } L^1((0, P) \times \mathbb{R})\end{aligned}$$

when $\varepsilon \rightarrow 0$.

Next, we pass to the limit in the symbol $a_i^\varepsilon(p, \xi)$:

Proposition 6.4. *The following limits hold pointwise $(p, \xi) \in (0, \infty) \times \mathbb{R}$ and strongly in $L^p_{loc}((0, \infty) \times \mathbb{R})$ for all $p \in (1, \infty)$:*

$$a_1^\varepsilon(p, \xi) \longrightarrow -p - \kappa_1 |\xi|^{\frac{8}{5}} \quad \text{with} \quad \kappa_1 = \frac{6}{5} \left(\frac{\pi}{v_0} \right)^{3/5} \int_0^\infty \frac{z^{3/5}}{z^2 + 1} dz \quad (54)$$

$$a_2^\varepsilon(p, \xi) \longrightarrow -\kappa_2 |\xi| \quad \text{with} \quad \kappa_2 = \frac{6}{5} \int_0^\infty \frac{1}{z^2 + 1} dz \quad (55)$$

$$a_3^\varepsilon(p, \xi) \longrightarrow -\kappa_3 |\xi|^{\frac{2}{5}} \quad \text{with} \quad \kappa_3 = \frac{6}{5} \left(\frac{v_0}{\pi} \right)^{3/5} \int_0^\infty \frac{z^{-3/5}}{z^2 + 1} dz \quad (56)$$

Furthermore, $a_1^\varepsilon, a_2^\varepsilon, a_3^\varepsilon \in L^\infty_{loc}((0, \infty) \times \mathbb{R})$ uniformly with respect to ε .

Finally, we need to show that the remainder terms, involving $f^\varepsilon - \Pi(f^\varepsilon)$, go to zero:

Proposition 6.5. *For all $0 < a < P$ and $K > 0$, we have*

$$R_i^\varepsilon \rightarrow 0 \quad \text{in } L^2((a, P) \times (-K, K))$$

as $\varepsilon \rightarrow 0$ for $i = 1, 2$.

Proof of Theorem 4.1. We are now ready to prove Theorem 4.1. First, using Proposition 6.1, we see that up to a subsequence, $T^\varepsilon(t, x)$ converges weakly to $T(t, x)$ in $L^2((0, \tau) \times \mathbb{R})$ for all τ (the uniqueness of the limit will give the convergence of the whole sequence).

Next, for a given test function $\varphi(p, \xi)$ in $\mathcal{D}((0, \infty) \times \mathbb{R})$, we then have

$$\int_0^\infty \int_{\mathbb{R}} \widehat{T}^\varepsilon(p, \xi) \varphi(p, \xi) d\xi dp = \int_0^\infty \int_{\mathbb{R}} T^\varepsilon(t, x) \widehat{\varphi}(t, x) dx dt \quad (57)$$

where $\widehat{\varphi} \in L^2((0, \infty) \times \mathbb{R})$. This last fact is the classical Parseval inequality for the Fourier transform, while for the Laplace transform, it follows from Minkowski's integral inequality:

$$\begin{aligned} \left(\int_0^\infty \left(\int_0^\infty e^{-pt} \varphi(p) dp \right)^2 dt \right)^{1/2} &\leq \int_0^\infty \left(\int_0^\infty e^{-2pt} dt \right)^{1/2} \varphi(p) dp \\ &\leq \int_0^\infty \frac{1}{\sqrt{2p}} \varphi(p) dp < \infty. \end{aligned}$$

Thus \widehat{T}^ε converges to \widehat{T} in $\mathcal{D}'((0, \infty) \times \mathbb{R})$. Since \widehat{T}^ε is also bounded in $L^2_{loc}((0, \infty) \times \mathbb{R})$ (using (48)), we deduce that (up to another subsequence) it converges weakly in $L^2_{loc}((0, \infty) \times \mathbb{R})$ to \widehat{T} .

In order to derive the equation satisfied by \widehat{T} , we need to pass to the limit in (52) and (53). However, we do not know that S^ε (defined in (46)) is bounded in some functional space. So we multiply equation (52) by a_3^ε and (53) by a_2^ε and consider their difference, in order to get rid of the terms in \widehat{S}^ε :

$$\begin{aligned} 0 &= a_3^\varepsilon(p, \xi) \mathcal{F}_1^\varepsilon(\widehat{f}^0) + \left(a_3^\varepsilon(p, \xi) a_1^\varepsilon(p, \xi) - (a_2^\varepsilon(p, \xi))^2 \right) \widehat{T}^\varepsilon(p, \xi) \\ &\quad + a_3^\varepsilon(p, \xi) R_1^\varepsilon(p, \xi) - a_2^\varepsilon(p, \xi) \mathcal{F}_2^\varepsilon(\widehat{f}^0) - a_2^\varepsilon(p, \xi) R_2^\varepsilon(p, \xi). \end{aligned}$$

Using Proposition 6.4, Proposition 6.5 and Proposition 6.3, we can now pass to the limit in this equation in $\mathcal{D}'((0, \infty) \times \mathbb{R})$ and deduce:

$$-\kappa_3 |\xi|^{2/5} \widehat{T}_0 + \left(-\kappa_3 |\xi|^{2/5} (-p - \kappa_1 |\xi|^{8/5}) - \kappa_2^2 |\xi|^2 \right) \widehat{T} = 0 \quad \text{in } \mathcal{D}'((0, \infty) \times \mathbb{R}).$$

Furthermore, factorizing $-\kappa_3 |\xi|^{2/5}$ in this last equation we get

$$-\kappa_3 |\xi|^{2/5} \left(\widehat{T}_0 - p \widehat{T} - \left(\kappa_1 + \frac{\kappa_2^2}{\kappa_3} \right) |\xi|^{8/5} \widehat{T} \right) = 0 \quad \text{in } \mathcal{D}'((0, \infty) \times \mathbb{R}).$$

This implies that the function

$$g(p, \xi) := \widehat{T}_0 - p \widehat{T} - \left(\kappa_1 + \frac{\kappa_2^2}{\kappa_3} \right) |\xi|^{8/5} \widehat{T}, \quad (58)$$

which belongs to $L_{loc}^2((0, \infty) \times \mathbb{R})$, satisfies

$$g(p, \xi) = 0 \text{ a.e. in } (0, \infty) \times \mathbb{R}$$

which gives (27)-(28).

To complete the proof of Theorem 4.1, it remains to show that f^ε converges to $T(t, x)$ (weakly in $L^\infty((0, \infty), L^2(\mathbb{R} \times \mathbb{T}))$). Since f^ε is bounded in $L^\infty(0, \infty; L^2(\mathbb{R} \times \mathbb{T}))$, and in view of the expansion (47), it is enough to show that $\varepsilon^{3/5} S^\varepsilon$ converges to zero in some weak sense.

This follows from Proposition 4.2, the proof of which uses equation (53) and some bounds from below on $a_3^\varepsilon(p, \xi)$ and will be detailed in Section 7. \square

We end this section by proving that the diffusion coefficient κ is indeed positive:

Lemma 6.6. *The coefficients κ_1 , κ_2 and κ_3 are such that*

$$\kappa_1 - \frac{\kappa_2^2}{\kappa_3} > 0.$$

Proof. Indeed, this is equivalent to

$$\kappa_2^2 < \kappa_1 \kappa_3$$

and using the explicit formula for κ_1 , κ_2 and κ_3 , we see that this is equivalent to

$$\left(\int_0^\infty \frac{1}{1+z^2} dz \right)^2 < \int_0^\infty \frac{z^{3/5}}{1+z^2} dz \int_0^\infty \frac{z^{-3/5}}{1+z^2} dz$$

which is an immediate consequence of Hölder inequality. \square

6.3 Proofs of the asymptotic results

We recall here that \mathbb{T} denotes the torus \mathbb{R}/\mathbb{Z} and that $\omega(k) = |\sin(\pi k)|$. Since the dispersion relation ω is degenerate at $k = 0 \pm n$, it will be easier in the computation below to work with k in the symmetric interval $(-\frac{1}{2}, \frac{1}{2})$ (when working with the interval $(0, 1)$, we have to deal with both endpoints 0 and 1). Note that the function ω is even in that interval and that

$$\omega'(k) = \operatorname{sgn}(k)\pi \cos(\pi k).$$

Finally, Proposition 5.2 implies:

Proposition 6.7. *The function $k \mapsto V(k)$ is even and non-negative on the interval $(-\frac{1}{2}, \frac{1}{2})$. Furthermore the function $W(k) := V(k)|k|^{-5/3}$ for $k \in (-\frac{1}{2}, \frac{1}{2})$ satisfies*

$$\lim_{k \rightarrow 0} W(k) = w_0 := v_0 \pi^{5/3}$$

and

$$C_0^{-1} \leq W(k) \leq C_0$$

for some $C_0 > 0$.

Proof of Proposition 6.3. The first part of the proposition follows immediately from Lebesgue dominated convergence theorem, since

$$\left| \frac{V(k)}{\varepsilon^{\frac{8}{5}} p + V(k) + i\varepsilon \omega'(k)\xi} \right| = \frac{V(k)}{\sqrt{\left(\varepsilon^{\frac{8}{5}} p + V(k)\right)^2 + (\varepsilon \omega'(k)\xi)^2}} \leq 1$$

and

$$\frac{V(k)}{\varepsilon^{\frac{8}{5}} p + V(k) + i\varepsilon \omega'(k)\xi} \rightarrow 1 \quad \text{as } \varepsilon \rightarrow 0.$$

For the second part, we note that

$$\left| \frac{V(k)}{\varepsilon^{\frac{8}{5}}p + V(k) + i\varepsilon\omega'(k)\xi} \right| \leq \frac{V(k)}{\varepsilon^{\frac{8}{5}}p + V(k)}$$

and so

$$\begin{aligned} |\mathcal{F}_2^\varepsilon(\widehat{f}^0)|(\xi, p) &\leq \varepsilon^{\frac{3}{5}} \int_{\mathbb{T}} \frac{V(k)}{\varepsilon^{\frac{8}{5}}p + V(k)} \frac{\widehat{f}_0(\xi, k)}{\omega(k)} dk \\ &\leq C\varepsilon^{\frac{3}{5}} \|\widehat{f}_0(\xi, \cdot)\|_{L^\infty(\mathbb{T})} \int_0^{1/2} \frac{|k|^{2/3}}{\varepsilon^{\frac{8}{5}}p + |k|^{5/3}} dk \\ &\leq C\varepsilon^{\frac{3}{5}} \|\widehat{f}_0(\xi, \cdot)\|_{L^\infty(\mathbb{T})} (1 + |\ln(\varepsilon^{\frac{8}{5}}p)|) \end{aligned} \quad (59)$$

hence the result, since this last inequality implies (integrating with respect to ξ and p)

$$\|\mathcal{F}_2^\varepsilon(\widehat{f}^0)\|_{L^1((0,P)\times\mathbb{R})} \leq C\varepsilon^{\frac{3}{5}} \|f_0\|_{L^\infty(\mathbb{R}\times\mathbb{T})} P(1 + |\ln(\varepsilon^{\frac{8}{5}}P)|).$$

□

Proof of Proposition 6.4. First, we write

$$\begin{aligned} 1 - \frac{V(k)}{\varepsilon^{\frac{8}{5}}p + V(k) + i\varepsilon\omega'(k)\xi} &= \frac{\varepsilon^{\frac{8}{5}}p + i\varepsilon\omega'(k)\xi}{\varepsilon^{\frac{8}{5}}p + V(k) + i\varepsilon\omega'(k)\xi} \\ &= \frac{\varepsilon^{\frac{8}{5}}p + V(k)}{(\varepsilon^{\frac{8}{5}}p + V(k))^2 + (\varepsilon\omega'(k)\xi)^2} \varepsilon^{\frac{8}{5}}p \\ &\quad + \frac{Vi\varepsilon\omega'(k)\xi}{(\varepsilon^{\frac{8}{5}}p + V(k))^2 + (\varepsilon\omega'(k)\xi)^2} \\ &\quad + \frac{(\varepsilon\omega'(k)\xi)^2}{(\varepsilon^{\frac{8}{5}}p + V(k))^2 + (\varepsilon\omega'(k)\xi)^2} \end{aligned} \quad (60)$$

Using the fact that $V(k) = V(-k)$, $\omega'(-k) = -\omega'(k)$, we deduce that

$$\begin{aligned} a_1^\varepsilon(p, \xi) &:= -p \int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{\varepsilon^{\frac{8}{5}}p + V(k)}{(\varepsilon^{\frac{8}{5}}p + V(k))^2 + (\varepsilon\omega'(k)\xi)^2} V(k) dk \\ &\quad - \varepsilon^{-\frac{8}{5}} \int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{(\varepsilon\omega'(k)\xi)^2}{(\varepsilon^{\frac{8}{5}}p + V(k))^2 + (\varepsilon\omega'(k)\xi)^2} V(k) dk. \end{aligned}$$

Dominated convergence immediately implies that the first term converges to $-p$, so we only have to consider the term

$$d^\varepsilon(p, \xi) = \varepsilon^{-\frac{8}{5}} \int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{(\varepsilon\omega'(k)\xi)^2}{(\varepsilon^{\frac{8}{5}}p + V(k))^2 + (\varepsilon\omega'(k)\xi)^2} V(k) dk.$$

For some $\delta \in (0, \frac{1}{2})$, we write

$$d^\varepsilon(p, \xi) = d_1^\varepsilon(p, \xi) + d_2^\varepsilon(p, \xi)$$

where

$$\begin{aligned} d_1^\varepsilon(p, \xi) &= \varepsilon^{-\frac{8}{5}} \int_{\substack{k \in (-\frac{1}{2}, \frac{1}{2}) \\ |k| \geq \delta}} \frac{(\varepsilon\omega'(k)\xi)^2}{(\varepsilon^{\frac{8}{5}}p + V(k))^2 + (\varepsilon\omega'(k)\xi)^2} V(k) dk \\ &\leq C\varepsilon^{-\frac{8}{5}} \int_{\substack{k \in (-\frac{1}{2}, \frac{1}{2}) \\ |k| \geq \delta}} \frac{(\varepsilon\xi)^2}{V(k)} dk \\ &\leq C(\delta)|\xi|^2 \varepsilon^{\frac{2}{5}} \end{aligned}$$

and

$$\begin{aligned} d_2^\varepsilon(p, \xi) &= \varepsilon^{-\frac{8}{5}} \int_{|k| \leq \delta} \frac{(\varepsilon\omega'(k)\xi)^2}{(\varepsilon^{\frac{8}{5}}p + V(k))^2 + (\varepsilon\omega'(k)\xi)^2} V(k) dk \\ &= 2\varepsilon^{-\frac{8}{5}} \int_0^\delta \frac{(\varepsilon\pi \cos(\pi k)\xi)^2}{(\varepsilon^{\frac{8}{5}}p + W(k)|k|^{5/3})^2 + (\varepsilon\pi \cos(\pi k)\xi)^2} W(k)|k|^{5/3} dk \\ &= 2\varepsilon^{-\frac{8}{5}} \int_0^\delta \frac{(\pi \cos(\pi k))^2}{(\varepsilon^{\frac{3}{5}}\frac{p}{|\xi|} + W(k)\frac{|k|^{5/3}}{\varepsilon|\xi|})^2 + (\pi \cos(\pi k))^2} W(k)|k|^{5/3} dk. \end{aligned}$$

We now do the change of variable

$$w = \frac{|k|^{5/3}}{\varepsilon|\xi|}, \quad dk = \frac{3}{5}(\varepsilon|\xi|)^{3/5} w^{-2/5} dw, \quad (61)$$

which yields

$$\begin{aligned} d_2^\varepsilon(p, \xi) &= 2\varepsilon^{-\frac{8}{5}} \int_0^{\frac{\delta^{5/3}}{\varepsilon|\xi|}} \frac{z^\varepsilon(w)}{\left(\varepsilon^{\frac{3}{5}}\frac{p}{|\xi|} + W^\varepsilon(w)\right)^2 + z^\varepsilon(w)} W^\varepsilon(w)\varepsilon|\xi|w\frac{3}{5}(\varepsilon|\xi|)^{3/5}w^{-2/5} dw \\ &= |\xi|^{8/5} \frac{6}{5} \int_0^{\frac{\delta^{5/3}}{\varepsilon|\xi|}} \frac{z^\varepsilon(w)}{\left(\varepsilon^{\frac{3}{5}}\frac{p}{|\xi|} + W^\varepsilon(w)\right)^2 + z^\varepsilon(w)} W^\varepsilon(w)w^{3/5} dw \end{aligned}$$

where

$$\begin{aligned} z^\varepsilon(w) &= (\pi \cos(\pi(\varepsilon|\xi|w)^{3/5}))^2 \\ W^\varepsilon(w) &= W((\varepsilon|\xi|w)^{3/5}). \end{aligned}$$

In particular, the integrand converges pointwise (for all w and ξ), as ε goes to zero, to

$$\frac{\pi^2}{(w_0 w)^2 + \pi^2} w_0 w^{3/5}$$

and it is bounded by

$$\frac{\pi^2}{(C_0^{-1} w)^2 + (\pi \cos(\pi \delta))^2} C_0 w^{3/5}.$$

We deduce that

$$|d_2^\varepsilon(p, \xi)| \leq C |\xi|^{8/5}$$

for some constant C and that

$$d_2^\varepsilon(p, \xi) \longrightarrow |\xi|^{8/5} \frac{6}{5} \int_0^\infty \frac{\pi^2}{(w_0 w)^2 + \pi^2} w_0 w^{3/5} dw = \kappa_1 |\xi|^{8/5}$$

(recall that $w_0 = v_0 \pi^{5/3}$) which concludes the proof of the first part. Note that we have also proved that

$$|a_1^\varepsilon(p, \xi)| \leq p + C \varepsilon^{2/5} |\xi|^2 + C |\xi|^{8/5}.$$

In particular, $a_1^\varepsilon(p, \xi)$ is bounded in $L_{loc}^\infty((0, \infty) \times \mathbb{R})$. Since it converges pointwise, a classical argument shows that it also converges strongly in $L_{loc}^p((0, \infty) \times \mathbb{R})$ for all $0 < p < \infty$.

The convergence of a_2^ε is proved similarly: Using (60), we find

$$\begin{aligned} a_2^\varepsilon(p, \xi) &:= -\varepsilon^{3/5} p \int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{\varepsilon^{8/5} p + V(k)}{(\varepsilon^{8/5} p + V(k))^2 + (\varepsilon \omega'(k) \xi)^2} \frac{V(k)}{\omega(k)} dk \\ &\quad - \varepsilon^{-1} \int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{(\varepsilon \omega'(k) \xi)^2}{(\varepsilon^{8/5} p + V(k))^2 + (\varepsilon \omega'(k) \xi)^2} \frac{V(k)}{\omega(k)} dk. \end{aligned}$$

The first term is bounded by

$$\begin{aligned}
& \varepsilon^{\frac{3}{5}} p \int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{1}{\varepsilon^{\frac{8}{5}} p + V(k)} \frac{V(k)}{\omega(k)} dk \\
& \leq C \varepsilon^{\frac{3}{5}} p \int_0^{\frac{1}{2}} \frac{1}{\varepsilon^{\frac{8}{5}} p + C_0^{-1} k^{\frac{5}{3}}} k^{\frac{2}{3}} dk \\
& \leq C \varepsilon^{-1} \int_0^{\frac{1}{2}} \frac{1}{1 + \varepsilon^{-\frac{8}{5}} p^{-1} k^{\frac{5}{3}}} k^{\frac{2}{3}} dk \\
& \leq C \varepsilon^{-1} p \varepsilon^{\frac{8}{5}} \int_0^{C \varepsilon^{-\frac{8}{5}} p^{-1}} \frac{1}{1+w} dw \\
& \leq C p \varepsilon^{\frac{3}{5}} \ln(1 + C \varepsilon^{-\frac{8}{5}} p^{-1})
\end{aligned}$$

and thus converges to zero as $\varepsilon \rightarrow 0$ (here we used the change of variable $w = \varepsilon^{-\frac{8}{5}} p^{-1} k^{\frac{5}{3}}$). For the second term the same decomposition of the integral in the interval $|k| \leq \delta$ and $|k| \geq \delta$. The integral in $|k| \geq \delta$ is bounded by $C(\delta)\varepsilon|\xi|^2$. For the integral in $|k| \leq \delta$, the change of variable (61) gives that it is bounded by $C|\xi|$ and converges to

$$|\xi| \frac{6}{5} \int_0^\infty \frac{\pi^2}{(w_0 w)^2 + \pi^2} \frac{w_0}{\pi} dw = \kappa_2 |\xi|.$$

Note that

$$|a_2^\varepsilon(p, \xi)| \leq C p \varepsilon^{\frac{3}{5}} \ln(1 + C \varepsilon^{-\frac{8}{5}} p^{-1}) + C(\delta)\varepsilon|\xi|^2 + C|\xi| \quad (62)$$

so $a_2^\varepsilon \in L_{loc}^\infty((0, \infty) \times \mathbb{R})$ implying, next to the pointwise convergence, the $L_{loc}^p((0, \infty) \times \mathbb{R})$ strong convergence for $0 < p < \infty$.

Finally, using (60), we find

$$\begin{aligned}
a_3^\varepsilon(p, \xi) & := -\varepsilon^{\frac{6}{5}} p \int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{\varepsilon^{\frac{8}{5}} p + V(k)}{(\varepsilon^{\frac{8}{5}} p + V(k))^2 + (\varepsilon \omega'(k) \xi)^2} \frac{V(k)}{\omega(k)^2} dk \\
& \quad - \varepsilon^{-\frac{2}{5}} \int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{(\varepsilon \omega'(k) \xi)^2}{(\varepsilon^{\frac{8}{5}} p + V(k))^2 + (\varepsilon \omega'(k) \xi)^2} \frac{V(k)}{\omega(k)^2} dk. \quad (63)
\end{aligned}$$

The first term is bounded by

$$\begin{aligned}
& \varepsilon^{\frac{6}{5}} p \int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{1}{\varepsilon^{\frac{8}{5}} p + V(k)} \frac{V(k)}{(\omega(k))^2} dk \\
& \leq C \varepsilon^{\frac{6}{5}} p \int_0^{\frac{1}{2}} \frac{1}{\varepsilon^{\frac{8}{5}} p + C_0^{-1} k^{\frac{5}{3}}} k^{-\frac{1}{3}} dk \\
& \leq C \varepsilon^{-\frac{2}{5}} \int_0^{\frac{1}{2}} \frac{1}{1 + \varepsilon^{-\frac{8}{5}} p^{-1} k^{\frac{5}{3}}} k^{-\frac{1}{3}} dk \\
& \leq C \varepsilon^{\frac{24}{25}} p^{\frac{4}{5}} \int_0^{C \varepsilon^{-\frac{8}{5}} p^{-1}} \frac{w^{-\frac{3}{5}}}{1+w} dw \\
& \leq C \varepsilon^{\frac{24}{25}} p^{\frac{4}{5}} \int_0^{\infty} \frac{w^{-\frac{3}{5}}}{1+w} dw
\end{aligned}$$

and thus converges to zero as $\varepsilon \rightarrow 0$. For the second term, the same decomposition of the integral in the interval $|k| \leq \delta$ and $|k| \geq \delta$. The integral in $|k| \geq \delta$ is bounded by $C(\delta)\varepsilon^{8/5}|\xi|$. The integral in $|k| \leq \delta$, the change of variable (61) gives that it is bounded by $C|\xi|^{2/5}$ and converges to

$$|\xi|^{\frac{2}{5}} \frac{6}{5} \int_0^{\infty} \frac{\pi^2}{(w_0 w)^2 + \pi^2} \frac{w_0}{\pi^2} w^{-\frac{3}{5}} dw = \kappa_3 |\xi|^{\frac{2}{5}}.$$

Analogously as in the previous cases, we have that

$$|a_3^\varepsilon(p, \xi)| \leq C \varepsilon^{\frac{24}{25}} p^{\frac{4}{5}} + C(\delta)\varepsilon^{8/5}|\xi| + C|\xi|^{2/5}$$

so $a_3^\varepsilon \in L_{loc}^\infty((0, \infty) \times \mathbb{R})$ which, next to the pointwise convergence, implies the $L_{loc}^p((0, \infty) \times \mathbb{R})$ strong convergence $p \in (0, \infty)$. \square

It only remain to prove Proposition 6.5. For that we will require the following lemma:

Lemma 6.8. *For all $\eta \in (0, \frac{1}{3}]$, we have*

$$\begin{aligned}
& \int_{\mathbb{T}} \left| \frac{V(k)}{\varepsilon^{\frac{8}{5}} p + V(k) + i\varepsilon\omega'(k)\xi} - 1 \right|^2 V(k) (\sin(\pi k))^{\frac{1}{3}-\eta} dk \\
& \leq C \left[(\varepsilon^{\frac{8}{5}} p)^{\frac{8}{5}} + (\varepsilon|\xi|)^{\frac{9}{5}-\frac{3\eta}{5}} + (\varepsilon|\xi|)^2 \right] \quad (64)
\end{aligned}$$

and

$$\int_{\mathbb{T}} \left| \frac{V(k)}{\varepsilon^{\frac{8}{5}}p + V(k) + i\varepsilon\omega'(k)\xi} - 1 \right|^2 \frac{V(k)}{\omega(k)^2} (\sin(\pi k))^{\frac{1}{3}-\eta} dk \leq C \left[(\varepsilon^{\frac{8}{5}}p)^{\frac{3}{5}(1-\eta)} + (\varepsilon\xi)^{\frac{3}{5}(1-\eta)} + (\varepsilon|\xi|)^2 \right] \quad (65)$$

We note that when $\eta = \frac{1}{3}$ (that is when we do not have the term $(\sin(\pi k))^{\frac{1}{3}-\eta}$ in the integral), then the integral behaves like $\varepsilon^{\frac{8}{5}}$. As we will see below, this would be just enough to show that the remainder term R_1^ε is bounded, but not to show that it converges to zero. The improvement of the norm of K given by (40) is thus essential here.

We first prove Proposition 6.5 (using Lemma 6.8), before giving the proof of Lemma 6.8:

Proof of Proposition 6.5. Using (40), we get:

$$\begin{aligned} |R_1^\varepsilon(p, \xi)| &= \varepsilon^{-\frac{8}{5}} \left| \int_{\mathbb{T}} \left(\frac{V(k)}{\varepsilon^{\frac{8}{5}}p + V(k) + i\varepsilon\omega'(k)\xi} - 1 \right) K(\widehat{f^\varepsilon} - \Pi(\widehat{f^\varepsilon}))(k) dk \right| \\ &\leq \varepsilon^{-\frac{8}{5}} \left(\int_{\mathbb{T}} \left| \frac{V(k)}{\varepsilon^{\frac{8}{5}}p + V(k) + i\varepsilon\omega'(k)\xi} - 1 \right|^2 V(k) (\sin(\pi k))^{\frac{1}{3}-\eta} dk \right)^{1/2} \\ &\quad \times \left(\int_{\mathbb{T}} K(\widehat{f^\varepsilon} - \Pi(\widehat{f^\varepsilon}))^2 (\sin(\pi k))^{-\frac{1}{3}+\eta} V^{-1}(k) dk \right)^{1/2} \\ &\leq C\varepsilon^{-\frac{8}{5}} \left(\int_{\mathbb{T}} \left| \frac{V(k)}{\varepsilon^{\frac{8}{5}}p + V(k) + i\varepsilon\omega'(k)\xi} - 1 \right|^2 V(k) (\sin(\pi k))^{\frac{1}{3}-\eta} dk \right)^{1/2} \\ &\quad \times \left(\int_{\mathbb{T}} (\widehat{f^\varepsilon} - \Pi(\widehat{f^\varepsilon}))^2 (\sin(\pi k))^{\frac{1}{3}-\eta} V(k) dk \right)^{1/2} \end{aligned}$$

and using (64), we deduce that for $p < P$ and $|\xi| \leq K$, we have

$$|R_1^\varepsilon(p, \xi)| \leq C(P, K) \varepsilon^{-\frac{8}{5}} \varepsilon^{\frac{9-3\eta}{10}} \left(\int_{\mathbb{T}} (\widehat{f^\varepsilon} - \Pi(\widehat{f^\varepsilon}))^2 V(k) dk \right)^{1/2}.$$

For all $0 < a < P$ and $K > 0$, we deduce

$$\begin{aligned}
\int_a^P \int_{-K}^K |R_1^\varepsilon(p, \xi)|^2 d\xi dp &\leq C(P, K) \varepsilon^{-\frac{16}{5}} \varepsilon^{\frac{9-3\eta}{5}} \|\widehat{f^\varepsilon} - \Pi(\widehat{f^\varepsilon})\|_{L^\infty((a, \infty); L_V^2(\mathbb{T} \times \mathbb{R}))}^2 \\
&\leq C(P, K) \frac{1}{2a} \varepsilon^{-\frac{7-3\eta}{5}} \|f^\varepsilon - \Pi(f^\varepsilon)\|_{L^2((0, \infty); L_V^2(\mathbb{T} \times \mathbb{R}))}^2 \\
&\leq C(a, P, K) \varepsilon^{-\frac{7-3\eta}{5}} \varepsilon^{\frac{8}{5}} \\
&\leq C(a, P, K) \varepsilon^{\frac{1-3\eta}{5}}
\end{aligned}$$

where we have used (48).

Clearly, this implies Proposition 6.5 for $i = 1$.

Proceeding similarly, we have that

$$\begin{aligned}
|R_2^\varepsilon(p, \xi)| &\leq C \varepsilon^{-1} \left((\varepsilon^{\frac{8}{5}} p)^{\frac{3}{5}(1-\eta)} + (\varepsilon \xi)^{\frac{3}{5}(1-\eta)} + (\varepsilon |\xi|)^2 \right)^{1/2} \\
&\quad \times \left(\int_{\mathbb{T}} (\widehat{f^\varepsilon} - \Pi(\widehat{f^\varepsilon}))^2 V(k) dk \right)^{1/2} \quad (66)
\end{aligned}$$

and therefore

$$\begin{aligned}
\int_a^P \int_{-K}^K |R_2^\varepsilon(p, \xi)|^2 dp d\xi &\leq C(P, K) \varepsilon^{-2} \varepsilon^{\frac{3(1-\eta)}{5}} \|\widehat{f^\varepsilon} - \Pi(\widehat{f^\varepsilon})\|_{L^\infty((a, \infty); L_V^2(\mathbb{T} \times \mathbb{R}))}^2 \\
&\leq C(a, P, K) \varepsilon^{-\frac{7-3\eta}{5}} \|f^\varepsilon - \Pi(f^\varepsilon)\|_{L^2((0, \infty); L_V^2(\mathbb{T} \times \mathbb{R}))}^2 \\
&\leq C(a, P, K) \varepsilon^{\frac{1-3\eta}{5}} \quad (67)
\end{aligned}$$

which converges to zero for any $\eta \in (0, \frac{1}{3})$. \square

Proof of Lemma 6.8. We write:

$$\begin{aligned}
&\int_{\mathbb{T}} \left| \frac{V(k)}{\varepsilon^{\frac{8}{5}} p + V(k) + i\varepsilon\omega'(k)\xi} - 1 \right|^2 V(k) (\sin(\pi k))^{\frac{1}{3}-\eta} dk \\
&= \int_{\mathbb{T}} \left| \frac{\varepsilon^{\frac{8}{5}} p + i\varepsilon\omega'(k)\xi}{\varepsilon^{\frac{8}{5}} p + V(k) + i\varepsilon\omega'(k)\xi} \right|^2 V(k) (\sin(\pi k))^{\frac{1}{3}-\eta} dk \\
&= \int_{\mathbb{T}} \frac{(\varepsilon^{\frac{8}{5}} p)^2 + (\varepsilon\omega'(k)\xi)^2}{(\varepsilon^{\frac{8}{5}} p + V(k))^2 + (\varepsilon\omega'(k)\xi)^2} V(k) (\sin(\pi k))^{\frac{1}{3}-\eta} dk \\
&= I_1 + I_2
\end{aligned}$$

where

$$\begin{aligned}
I_1 &:= \int_{\mathbb{T}} \frac{(\varepsilon^{\frac{8}{5}}p)^2}{(\varepsilon^{\frac{8}{5}}p + V(k))^2 + (\varepsilon\omega'(k)\xi)^2} V(k)(\sin(\pi k))^{\frac{1}{3}-\eta} dk \\
&\leq 2 \int_0^{1/2} \frac{(\varepsilon^{\frac{8}{5}}p)^2}{(\varepsilon^{\frac{8}{5}}p + V(k))^2} V(k)(\sin(\pi k))^{\frac{1}{3}-\eta} dk \\
&\leq 2 \int_0^{1/2} \frac{(\varepsilon^{\frac{8}{5}}p)^2}{(\varepsilon^{\frac{8}{5}}p + k^{5/3})^2} k^{5/3} dk
\end{aligned}$$

(note we do not need to use the $(\sin(\pi k))^{\frac{1}{3}-\eta}$ to control this term) and

$$\begin{aligned}
I_2 &:= \int_{\mathbb{T}} \frac{(\varepsilon\omega'(k)\xi)^2}{(\varepsilon^{\frac{8}{5}}p + V(k))^2 + (\varepsilon\omega'(k)\xi)^2} V(k)(\sin(\pi k))^{\frac{1}{3}-\eta} dk \\
&\leq 2 \int_0^{1/4} \frac{(\varepsilon\omega'(k)\xi)^2}{V(k)^2 + (\varepsilon\omega'(k)\xi)^2} V(k)(\sin(\pi k))^{\frac{1}{3}-\eta} dk + 2 \int_{1/4}^{1/2} \frac{(\varepsilon\pi\xi)^2}{V(k)} dk \\
&\leq C \int_0^{1/4} \frac{(\varepsilon\xi)^2}{k^{10/3} + (\varepsilon\xi)^2} k^{2-\eta} dk + C\varepsilon^2|\xi|^2
\end{aligned}$$

(here the $(\sin(\pi k))^{\frac{1}{3}-\eta}$ is essential).

Using the change of variable $w = \frac{k^{5/3}}{\varepsilon^{8/5}p}$ in I_1 and $w = \frac{k^{5/3}}{\varepsilon\xi}$ in I_2 , we find

$$\begin{aligned}
I_1 &\leq C(\varepsilon^{8/5}p)^{8/5} \int_0^\infty \frac{w^{3/5}}{(1+w)^2} dw \\
I_2 &\leq C(\varepsilon\xi)^{9/5-3\eta/5} \int_0^\infty \frac{w^{4/5-3\eta/5}}{1+w^2} dw + C\varepsilon^2|\xi|^2
\end{aligned}$$

where the integral in the right hand side are clearly finite (recall that $\eta \in (0, \frac{1}{3})$). Inequality (64) follows.

We now proceed similarly to prove (65): First, we write

$$\begin{aligned}
&\int_{\mathbb{T}} \left| \frac{V(k)}{\varepsilon^{\frac{8}{5}}p + V(k) + i\varepsilon\omega'(k)\xi} - 1 \right|^2 \frac{V(k)}{\omega(k)^2} (\sin(\pi k))^{\frac{1}{3}-\eta} dk \\
&= \int_{\mathbb{T}} \frac{(\varepsilon^{\frac{8}{5}}p)^2 + (\varepsilon\omega'(k)\xi)^2}{(\varepsilon^{\frac{8}{5}}p + V(k))^2 + (\varepsilon\omega'(k)\xi)^2} \frac{V(k)}{\omega(k)^2} (\sin(\pi k))^{\frac{1}{3}-\eta} dk \\
&= \tilde{I}_1 + \tilde{I}_2
\end{aligned}$$

where

$$\begin{aligned}
\tilde{I}_1 &:= \int_{\mathbb{T}} \frac{(\varepsilon^{\frac{8}{5}}p)^2}{(\varepsilon^{\frac{8}{5}}p + V(k))^2 + (\varepsilon\omega'(k)\xi)^2} \frac{V(k)}{\omega(k)^2} (\sin(\pi k))^{\frac{1}{3}-\eta} dk \\
&\leq 2 \int_0^{1/2} \frac{(\varepsilon^{\frac{8}{5}}p)^2}{(\varepsilon^{\frac{8}{5}}p + V(k))^2} \frac{V(k)}{\omega(k)^2} (\sin(\pi k))^{\frac{1}{3}-\eta} dk \\
&\leq 2 \int_0^{1/2} \frac{(\varepsilon^{\frac{8}{5}}p)^2}{(\varepsilon^{\frac{8}{5}}p + k^{5/3})^2} k^{-\eta} dk
\end{aligned}$$

and

$$\begin{aligned}
\tilde{I}_2 &:= \int_{\mathbb{T}} \frac{(\varepsilon\omega'(k)\xi)^2}{(\varepsilon^{\frac{8}{5}}p + V(k))^2 + (\varepsilon\omega'(k)\xi)^2} \frac{V(k)}{\omega(k)^2} (\sin(\pi k))^{\frac{1}{3}-\eta} dk \\
&\leq 2 \int_0^{1/2} \frac{(\varepsilon\omega'(k)\xi)^2}{V(k)^2 + (\varepsilon\omega'(k)\xi)^2} \frac{V(k)}{\omega(k)^2} (\sin(\pi k))^{\frac{1}{3}-\eta} dk \\
&\leq 2 \int_0^{1/4} \frac{(\varepsilon\xi)^2}{k^{10/3} + (\varepsilon\xi)^2} k^{-\eta} dk + C(\varepsilon|\xi|)^2
\end{aligned}$$

Using the change of variable $w = \frac{k^{5/3}}{\varepsilon^{8/5}p}$ in \tilde{I}_1 and $w = \frac{k^{5/3}}{\varepsilon\xi}$ in \tilde{I}_2 , we find

$$\begin{aligned}
\tilde{I}_1 &\leq C(\varepsilon^{\frac{8}{5}}p)^{\frac{3}{5}(1-\eta)} \int_0^\infty \frac{w^{-3/5\eta}}{(1+w)^2} dw \\
\tilde{I}_2 &\leq C(\varepsilon\xi)^{\frac{3}{5}(1-\eta)} \int_0^\infty \frac{w^{-2/5-3\eta/5}}{1+w^2} dw + C(\varepsilon|\xi|)^2
\end{aligned}$$

which yields (65). □

7 Proof of Proposition 4.2

The proof of Proposition 4.2 relies on the following crucial bound:

Lemma 7.1. *There exists a constant c such that for all K and for all ε such that $\varepsilon K \leq 1$, the following lower bound holds*

$$|a_3^\varepsilon(p, \xi)| \geq c\varepsilon^{\frac{6}{25}}p^{\frac{2}{5}} + c|\xi|^{\frac{2}{5}} \quad \text{for } 0 \leq p \leq K, \quad |\xi| \leq K. \quad (68)$$

Proof of Lemma 7.1. We recall that $a_3^\varepsilon(p, \xi)$ is given by (63). In particular, we note that for all $(p, \xi) \neq (0, 0)$, we have $a_3^\varepsilon(p, \xi) < 0$. Furthermore, we can write (using the fact that all the terms in (63) have the same sign):

$$\begin{aligned} -a_3^\varepsilon(p, k) &\geq \varepsilon^{\frac{6}{5}} p \int_0^{\frac{1}{4}} \frac{\varepsilon^{\frac{8}{5}} p + V(k)}{(\varepsilon^{\frac{8}{5}} p + V(k))^2 + (\varepsilon \omega'(k) \xi)^2} \frac{V(k)}{\omega(k)^2} dk \\ &\quad + \varepsilon^{-\frac{2}{5}} \int_0^{\frac{1}{4}} \frac{(\varepsilon \omega'(k) \xi)^2}{(\varepsilon^{\frac{8}{5}} p + V(k))^2 + (\varepsilon \omega'(k) \xi)^2} \frac{V(k)}{\omega(k)^2} dk. \end{aligned}$$

Using the fact that for $k \in (0, 1/4)$ we have $C_0^{-1}|k|^{5/3} \leq V(k) \leq C_0|k|^{5/3}$, $\frac{\pi}{\sqrt{2}} \leq \omega'(k) \leq \pi$ and $\frac{\pi}{2}k \leq \omega(k) \leq \pi k$, we obtain the following lower bound (for some constant $c > 0$):

$$\begin{aligned} -a_3^\varepsilon(p, k) &\geq c \varepsilon^{\frac{6}{5}} p \int_0^{\frac{1}{4}} \frac{|k|^{4/3}}{(\varepsilon^{\frac{8}{5}} p + C_0|k|^{5/3})^2 + (\varepsilon \pi \xi)^2} dk \\ &\quad + c \varepsilon^{-\frac{2}{5}} \int_0^{\frac{1}{4}} \frac{(\varepsilon \pi \xi)^2 k^{-1/3}}{(\varepsilon^{\frac{8}{5}} p + C_0|k|^{5/3})^2 + (\varepsilon \pi \xi)^2} dk. \end{aligned} \quad (69)$$

From now on, we fix K and assume that $0 < p \leq K$ and that $|\xi| \leq K$. We also assume that ε is such that $\varepsilon K \leq 1$. In order to establish (68), we consider two cases, and in each case we use only one of the integrals in (69):

1. First, assume that p and ξ are such that

$$|\xi| \leq \varepsilon^{\frac{3}{5}} p. \quad (70)$$

Then, using only the first integral in (69), we get (using (70)):

$$-a_3^\varepsilon(p, k) \geq c \varepsilon^{\frac{6}{5}} p \int_0^{\frac{1}{4}} \frac{|k|^{4/3}}{(\varepsilon^{\frac{8}{5}} p + C_0|k|^{5/3})^2 + (\pi \varepsilon^{\frac{8}{5}} p)^2} dk$$

and the change of variable $w = (\varepsilon^{\frac{8}{5}} p)^{-\frac{3}{5}} k$ yields

$$-a_3^\varepsilon(p, k) \geq c \varepsilon^{\frac{6}{5}} p \frac{(\varepsilon^{\frac{8}{5}} p)^{\frac{7}{5}}}{(\varepsilon^{\frac{8}{5}} p)^2} \int_0^{\frac{1}{4(\varepsilon^{\frac{8}{5}} p)^{\frac{3}{5}}}} \frac{|w|^{4/3}}{(1 + C_0|w|^{5/3})^2 + \pi^2} dw$$

and using the fact that $\varepsilon^{\frac{8}{5}} p \leq 1$, we deduce (for a different constant c):

$$-a_3^\varepsilon(p, k) \geq c \varepsilon^{\frac{6}{25}} p^{\frac{2}{5}}.$$

Finally, using (70), we also get

$$-a_3^\varepsilon(p, k) \geq c|\xi|^{\frac{2}{5}}$$

and so (68) holds in this case.

2. Next, we assume that p and ξ are such that

$$\varepsilon^{\frac{3}{5}}p \leq |\xi| \tag{71}$$

and using only the second integral in (69), we get (using (71)):

$$-a_3^\varepsilon(p, k) \geq c\varepsilon^{-\frac{2}{5}} \int_0^{\frac{1}{4}} \frac{(\varepsilon\pi\xi)^2 k^{-1/3}}{(\varepsilon|\xi| + C_0|k|^{5/3})^2 + (\varepsilon\pi\xi)^2} dk$$

and the change of variable $w = (\varepsilon|\xi|)^{-\frac{3}{5}}k$, yields:

$$\begin{aligned} -a_3^\varepsilon(p, k) &\geq c\varepsilon^{-\frac{2}{5}}\pi^2 \int_0^{\frac{1}{4(\varepsilon\xi)^{\frac{3}{5}}}} \frac{(\varepsilon\xi)^{\frac{2}{5}}w^{-1/3}}{(1 + C_0|w|^{5/3})^2 + \pi^2} dw \\ &\geq c|\xi|^{\frac{2}{5}} \end{aligned}$$

(using the fact that $\varepsilon|\xi| \leq 1$). Finally, using (71), we also get

$$-a_3^\varepsilon(p, k) \geq c\varepsilon^{\frac{6}{25}}p^{\frac{2}{5}}$$

and so (68) holds also in this case. □

Proof of Proposition 4.2. We use equation (53) to determine \widehat{S}^ε :

$$\widehat{S}^\varepsilon = \frac{1}{a_3^\varepsilon} \left(-\mathcal{F}_2^\varepsilon(\widehat{f}_0) - R_2^\varepsilon - a_2^\varepsilon \widehat{T}^\varepsilon \right). \tag{72}$$

Note that we can do this since $a_3^\varepsilon(p, \xi) < 0$ as long as p and ξ are not simultaneously zero.

We now need to show that we can pass to the limit in all the terms in the right hand side. First, using Lemma 7.1 and the estimate (62), we deduce that for a given K and for all $\varepsilon \leq K^{-1}$, we have

$$\begin{aligned} \frac{a_2^\varepsilon(p, \xi)}{a_3^\varepsilon(p, \xi)} &= \left| \frac{a_2^\varepsilon(p, \xi)}{a_3^\varepsilon(p, \xi)} \right| \leq \frac{Cp\varepsilon^{3/5} \ln(1 + C\varepsilon^{-8/5}p^{-1})}{c\varepsilon^{\frac{6}{25}}p^{\frac{2}{5}}} + \frac{C\varepsilon|\xi|^2}{c|\xi|^{\frac{2}{5}}} + \frac{C|\xi|}{c|\xi|^{\frac{2}{5}}} \\ &\leq Cp^{\frac{3}{5}}\varepsilon^{\frac{9}{25}} \ln(1 + C\varepsilon^{-8/5}p^{-1}) + C\varepsilon|\xi|^{\frac{8}{5}} + C|\xi|^{\frac{3}{5}} \\ &\leq C(K) \end{aligned}$$

for all $0 \leq p \leq K$ and $|\xi| \leq K$. Furthermore, this uniform bound, together with Proposition 6.4 implies that

$$\frac{a_2^\varepsilon(p, \xi)}{a_3^\varepsilon(p, \xi)} \longrightarrow \frac{\kappa_2}{\kappa_3} |\xi|^{3/5}$$

point-wise and in $L_{loc}^p((0, \infty) \times \mathbb{R})$ strong.

Next, for ε sufficiently small we can use Lemma 7.1 along with the estimates on $\mathcal{F}_2^\varepsilon(\hat{f}_0)$ in (59) to conclude that

$$\frac{1}{a_3^\varepsilon} \mathcal{F}_2^\varepsilon(\hat{f}_0) \rightarrow 0 \quad \text{in } \mathcal{D}'((0, \infty) \times \mathbb{R}).$$

Finally, we need to bound the quantity

$$\left| \frac{R_2^\varepsilon(p, \xi)}{a_3^\varepsilon(p, \xi)} \right|.$$

For that, we fix $0 < a < P$ and for $p \in (a, P)$ and $|\xi| \leq K$, estimate (66) then implies

$$\begin{aligned} |R_2^\varepsilon(p, \xi)| &\leq C\varepsilon^{-1} \left(\varepsilon^{\frac{24}{25}(1-\eta)} + (\varepsilon\xi)^{\frac{3}{5}(1-\eta)} \right)^{1/2} \left(\int_{\mathbb{T}} (\hat{f}^\varepsilon - \Pi(\hat{f}^\varepsilon))^2 V(k) dk \right)^{1/2} \\ &\leq \begin{cases} C\varepsilon^{-1} \varepsilon^{\frac{12}{25}(1-\eta)} \left(\int_{\mathbb{T}} (\hat{f}^\varepsilon - \Pi(\hat{f}^\varepsilon))^2 V(k) dk \right)^{1/2} & \text{if } |\xi| \leq \varepsilon^{3/5} \\ C\varepsilon^{-1} (\varepsilon\xi)^{\frac{3}{10}(1-\eta)} \left(\int_{\mathbb{T}} (\hat{f}^\varepsilon - \Pi(\hat{f}^\varepsilon))^2 V(k) dk \right)^{1/2} & \text{if } |\xi| \geq \varepsilon^{3/5} \end{cases} \end{aligned}$$

and we are going to use the following consequence of Lemma 7.1:

$$-a_3^\varepsilon(p, \xi) \geq \begin{cases} c(a)\varepsilon^{6/25} & \text{if } |\xi| \leq \varepsilon^{3/5} \\ c|\xi|^{2/5} & \text{if } |\xi| \geq \varepsilon^{3/5} \end{cases}$$

We deduce

$$\left| \frac{R_2^\varepsilon(p, \xi)}{a_3^\varepsilon(p, \xi)} \right| \leq \begin{cases} C\varepsilon^{\frac{-19-12\eta}{25}} \left(\int_{\mathbb{T}} (\hat{f}^\varepsilon - \Pi(\hat{f}^\varepsilon))^2 V(k) dk \right)^{1/2} & \text{if } |\xi| \leq \varepsilon^{3/5} \\ C\varepsilon^{-1} \varepsilon^{\frac{3}{10}(1-\eta)} |\xi|^{\frac{-1-3\eta}{10}} \left(\int_{\mathbb{T}} (\hat{f}^\varepsilon - \Pi(\hat{f}^\varepsilon))^2 V(k) dk \right)^{1/2} & \text{if } |\xi| \geq \varepsilon^{3/5} \end{cases}.$$

Finally, using the condition $|\xi| \geq \varepsilon^{3/5}$ in the second case, we deduce that

$$\left| \frac{R_2^\varepsilon(p, \xi)}{a_3^\varepsilon(p, \xi)} \right| \leq C(a, p, K) \varepsilon^{\frac{-19-12\eta}{25}} \left(\int_{\mathbb{T}} (\hat{f}^\varepsilon - \Pi(\hat{f}^\varepsilon))^2 V(k) dk \right)^{1/2}$$

for all $p \in (a, P)$ and $|\xi| \leq K$.

We deduce

$$\begin{aligned} \left(\int_a^P \int_{-K}^K \left| \frac{R_2^\varepsilon(p, \xi)}{a_3^\varepsilon(p, \xi)} \right|^2 d\xi dp \right)^{1/2} &\leq C(a, P, K) \varepsilon^{\frac{-19-12\eta}{25}} \|(\widehat{f^\varepsilon} - \Pi(\widehat{f^\varepsilon}))\|_{L^\infty((a, \infty); L_V^2(\mathbb{T} \times \mathbb{R}))} \\ &\leq C(a, P, K) \varepsilon^{\frac{-19-12\eta}{25}} \|f^\varepsilon - \Pi(f^\varepsilon)\|_{L^2((0, \infty); L_V^2(\mathbb{T} \times \mathbb{R}))} \\ &\leq C(a, P, K) \varepsilon^{\frac{1-12\eta}{25}} \end{aligned}$$

which goes to zero as $\varepsilon \rightarrow 0$.

We can now pass to the limit in (72) to conclude that

$$\widehat{S}^\varepsilon \longrightarrow \widehat{S} = \frac{\kappa_2}{\kappa_3} |\xi|^{3/5} \widehat{T} \quad \text{in } \mathcal{D}'((0, \infty) \times \mathbb{R})$$

which completes the proof of Proposition 4.2. □

References

- [1] N.B. Abdallah, A. Mellet, and M. Puel. Fractional diffusion limit for collisional kinetic equations: a hilbert expansion approach. *preprint*, 2010.
- [2] G Allaire and F Golse. Transport et diffusion. *Lecture notes, Ecole polytechnique (in French)*, 2013.
- [3] Kenichiro Aoki and Dimitri Kusnezov. Fermi-pasta-ulam β model: Boundary jumps, fourier's law, and scaling. *Phys. Rev. Lett.*, 86:4029–4032, Apr 2001.
- [4] Claude Bardos, François Golse, and David Levermore. Fluid dynamic limits of kinetic equations. i. formal derivations. *Journal of statistical physics*, 63(1-2):323–344, 1991.
- [5] Giada Basile, Cédric Bernardin, and Stefano Olla. Momentum conserving model with anomalous thermal conductivity in low dimensional systems. *Physical review letters*, 96(20):204303, 2006.
- [6] Giada Basile, Cédric Bernardin, and Stefano Olla. Thermal conductivity for a momentum conservative model. *Communications in Mathematical Physics*, 287(1):67–98, 2009.

- [7] Giada Basile, Stefano Olla, and Herbert Spohn. Energy transport in stochastically perturbed lattice dynamics. *Archive for rational mechanics and analysis*, 195(1):171–203, 2010.
- [8] Naoufel Ben Abdallah, Antoine Mellet, and Marjolaine Puel. Anomalous diffusion limit for kinetic equations with degenerate collision frequency. *Math. Models Methods Appl. Sci.*, 21(11):2249–2262, 2011.
- [9] GP Berman and FM Izrailev. The fermi–pasta–ulam problem: fifty years of progress. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 15(1):015104, 2005.
- [10] Federico Bonetto, Joel L Lebowitz, and Luc Rey-Bellet. Fourier’s law: A challenge for theorists. *arXiv preprint math-ph/0002052*, 2000.
- [11] Jean Bricmont and Antti Kupiainen. Approach to equilibrium for the phonon boltzmann equation. *Communications in Mathematical Physics*, 281(1):179–202, 2008.
- [12] Milton Jara, Tomasz Komorowski, and Stefano Olla. Limit theorems for additive functionals of a Markov chain. *Ann. Appl. Probab.*, 19(6):2270–2300, 2009.
- [13] Stefano Lepri, Roberto Livi, and Antonio Politi. Thermal conduction in classical low-dimensional lattices. *Physics Reports*, 377(1):1–80, 2003.
- [14] Stefano Lepri, Roberto Livi, and Antonio Politi. Studies of thermal conductivity in fermi–pasta–ulam-like lattices. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 15(1):015118, 2005.
- [15] Jani Lukkarinen and Herbert Spohn. Anomalous energy transport in the fpv- β chain. *Communications on Pure and Applied Mathematics*, 61(12):1753–1786, 2008.
- [16] Antoine Mellet. Fractional diffusion limit for collisional kinetic equations: a moments method. *Indiana Univ. Math. J.*, 59(4):1333–1360, 2010.
- [17] Antoine Mellet, Stéphane Mischler, and Clément Mouhot. Fractional diffusion limit for collisional kinetic equations. *Arch. Ration. Mech. Anal.*, 199(2):493–525, 2011.
- [18] Clément Mouhot. Lecture notes: Mathematical topics in kinetic theory. *Chapter 4: The linear Boltzmann equation*, 2013.

- [19] Stefano Olla. Energy diffusion and superdiffusion in oscillators lattice networks. In *New Trends in Mathematical Physics*, pages 539–547. Springer, 2009.
- [20] Rudolf Peierls. Zur kinetischen theorie der wärmeleitung in kristallen. *Annalen der Physik*, 395(8):1055–1101, 1929.
- [21] Andrey Pereverzev. Fermi-Pasta-Ulam β lattice: Peierls equation and anomalous heat conductivity. *Phys. Rev. E (3)*, 68(5):056124, 6, 2003.
- [22] Takashi Shimada, Teruyoshi Murakami, Satoshi Yukawa, Keiji Saito, and Nobuyasu Ito. Simulational study on dimensionality dependence of heat conduction. *Journal of the Physical Society of Japan*, 69(10):3150–3153, 2000.
- [23] Herbert Spohn. Collisional invariants for the phonon boltzmann equation. *Journal of statistical physics*, 124(5):1131–1135, 2006.
- [24] Herbert Spohn. The phonon boltzmann equation, properties and link to weakly anharmonic lattice dynamics. *Journal of statistical physics*, 124(2-4):1041–1104, 2006.
- [25] Herbert Spohn. On the Boltzmann equation for weakly nonlinear wave equations. In *Boltzmann’s legacy*, ESI Lect. Math. Phys., pages 145–159. Eur. Math. Soc., Zürich, 2008.

ANTOINE MELLET

UNIVERSITY OF MARYLAND
DEPARTMENT OF MATHEMATICS
COLLEGE PARK 20742
USA

E-MAIL: mellet@math.umd.edu

SARA MERINO-ACEITUNO

UNIVERSITY OF CAMBRIDGE
DPMMS, CENTRE FOR MATHEMATICAL SCIENCES
WILBERFORCE ROAD, CAMBRIDGE CB3 0WA, UK

E-MAIL: s.merino-aceituno@maths.cam.ac.uk